

This Page Is Inserted by IFW Operations
and is not a part of the Official Record

BEST AVAILABLE IMAGES

Defective images within this document are accurate representations of the original documents submitted by the applicant.

Defects in the images may include (but are not limited to):

- BLACK BORDERS
- TEXT CUT OFF AT TOP, BOTTOM OR SIDES
- FADED TEXT
- ILLEGIBLE TEXT
- SKEWED/SLANTED IMAGES
- COLORED PHOTOS
- BLACK OR VERY BLACK AND WHITE DARK PHOTOS
- GRAY SCALE DOCUMENTS

IMAGES ARE BEST AVAILABLE COPY.

**As rescanning documents *will not* correct images,
please do not report the images to the
Image Problem Mailbox.**

THIS PAGE BLANK (USF10,

PCTWORLD INTELLECTUAL PROPERTY ORGANIZATION
International Bureau

R0134A-REG

B6

INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ⁶ : C07D 209/16, A01N 43/38, A61K 31/40, 31/405	A1	(11) International Publication Number: WO 99/07677 (43) International Publication Date: 18 February 1999 (18.02.99)
(21) International Application Number: PCT/US98/16333 (22) International Filing Date: 6 August 1998 (06.08.98) (30) Priority Data: 60/054,997 6 August 1997 (06.08.97) US (71) Applicant (for all designated States except US): ELI LILLY AND COMPANY [US/US]; Lilly Corporate Center, Indianapolis, IN 46285 (US). (72) Inventors; and (75) Inventors/Applicants (for US only): FRITZ, James, Erwin [US/US]; 9757 North Moonstone Place, McCordsville, IN 46055 (US). HIPSKIND, Philip, Arthur [US/US]; 3660 South Farmstone Circle, New Palestine, IN 46163 (US). KALDOR, Stephen, Warren [US/US]; 7435 Burkwood Way, Indianapolis, IN 46254 (US). LOBB, Karen, Lynn [US/US]; 5625 East Lowell Avenue, Indianapolis, IN 46219 (US). NIXON, James, Arthur [US/US]; 7375 Taos Trail, Indianapolis, IN 46219 (US). (74) Agents: GAYLO, Paul, J. et al.; Eli Lilly and Company, Lilly Corporate Center, Indianapolis, IN 46285 (US).		(81) Designated States: AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CU, CZ, DE, DK, EE, ES, FI, GB, GE, GH, GM, HR, HU, ID, IL, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, UA, UG, US, UZ, VN, YU, ZW, ARIPO patent (GH, GM, KE, LS, MW, SD, SZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG). Published <i>With international search report.</i>
(54) Title: 2-ACYLAMINOPROPANAMINES AS TACHYKININ RECEPTOR ANTAGONISTS		
(57) Abstract Disclosed are non-peptidyl 2-acylaminopropanamine tachykinin receptor antagonists which are useful in the treatment of various conditions including Alzheimer's disease, and pharmaceutical compositions containing such compounds.		

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav Republic of Macedonia	TM	Turkmenistan
BF	Burkina Faso	GR	Greece	ML	Mali	TR	Turkey
BG	Bulgaria	HU	Hungary	MN	Mongolia	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MR	Mauritania	UA	Ukraine
BR	Brazil	IL	Israel	MW	Malawi	UG	Uganda
BY	Belarus	IS	Iceland	MX	Mexico	US	United States of America
CA	Canada	IT	Italy	NE	Niger	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NL	Netherlands	VN	Viet Nam
CG	Congo	KE	Kenya	NO	Norway	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NZ	New Zealand	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's Republic of Korea	PL	Poland		
CM	Cameroon	KR	Republic of Korea	PT	Portugal		
CN	China	KZ	Kazakhstan	RO	Romania		
CU	Cuba	LC	Saint Lucia	RU	Russian Federation		
CZ	Czech Republic	LI	Liechtenstein	SD	Sudan		
DE	Germany	LK	Sri Lanka	SE	Sweden		
DK	Denmark	LR	Liberia	SG	Singapore		
EE	Estonia						

- 1 -

Title**2-ACYLAMINOPROPANAMINES AS TACHYKININ RECEPTOR
ANTAGONISTS**

5

Background of the Invention

Tachykinins are a family of peptides which share a common amidated carboxy terminal sequence. Substance P was the first peptide of this family to be isolated, although its purification and the determination of its primary sequence did not occur until the early 1970's.

Between 1983 and 1984 several groups reported the isolation of two novel mammalian tachykinins, now termed neurokinin A (also known as substance K, neuromedin L, and neurokinin α), and neurokinin B (also known as neuromedin K and neurokinin β). See, J.E. Maggio, Peptides, 6 (Supplement 3):237-243 (1985) for a review of these discoveries.

Tachykinins are widely distributed in both the central and peripheral nervous systems, are released from nerves, and exert a variety of biological actions, which, in most cases, depend upon activation of specific receptors expressed on the membrane of target cells. Tachykinins are also produced by a number of non-neural tissues.

The mammalian tachykinins substance P, neurokinin A, and neurokinin B act through three major receptor subtypes, denoted as NK-1, NK-2, and NK-3, respectively. These receptors are present in a variety of organs.

Substance P is believed inter alia to be involved in the neurotransmission of pain sensations, including the pain associated with migraine headaches and with arthritis. These peptides have also been implicated in gastrointestinal disorders and diseases of the gastrointestinal tract such as inflammatory bowel disease. Tachykinins have also been implicated as playing a role in numerous other maladies, as discussed infra.

Tachykinins play a major role in mediating the sensation and transmission of pain or nociception, especially migraine headaches.

- 2 -

see, e.g., S.L. Shepheard, et al., British Journal of Pharmacology, 108:11-20 (1993); S.M. Moussaoui, et al., European Journal of Pharmacology, 238:421-424 (1993); and W.S. Lee, et al., British Journal of Pharmacology, 112:920-924 (1994).

5 In view of the wide number of clinical maladies associated with an excess of tachykinins, the development of tachykinin receptor antagonists will serve to control these clinical conditions. The earliest tachykinin receptor antagonists were peptide derivatives. These antagonists proved to be of limited pharmaceutical utility because of their
10 metabolic instability.

 Recent publications have described novel classes of non-peptidyl tachykinin receptor antagonists which generally have greater oral bioavailability and metabolic stability than the earlier classes of tachykinin receptor antagonists. Examples of such newer non-peptidyl
15 tachykinin receptor antagonists are found in United States Patent 5,491,140, issued February 13, 1996; United States Patent 5,328,927, issued July 12, 1994; United States Patent 5,360,820, issued November 1, 1994; United States Patent 5,344,830, issued September 6, 1994; United States Patent 5,331,089, issued July 19, 1994; European Patent
20 Publication 591,040 A1, published April 6, 1994; Patent Cooperation Treaty publication WO 94/01402, published January 20, 1994; Patent Cooperation Treaty publication WO 94/04494, published March 3, 1994; Patent Cooperation Treaty publication WO 93/011609, published January 21, 1993; Canadian Patent Application 2154116, published January 23,
25 1996; European Patent Publication 693,489, published January 24, 1996; and Canadian Patent Application 2151116, published December 11, 1995.

 United States Patent 5,530,009, issued June 25, 1996, describes a 1,2-diacylaminopropane for use in treating conditions associated with an excess of tachykinins. This patent also teaches
30 processes for preparing this compound.

 In essence, this invention provides a class of potent non-peptidyl tachykinin receptor antagonists similar to those of United States Patent 5,530,009. By virtue of their non-peptidyl nature, the compounds of the present invention do not suffer from the shortcomings, in terms of


- 3 -

metabolic instability, of known peptide-based tachykinin receptor antagonists.

Summary of the Invention

5

This invention provides novel compounds of Formula I



10

wherein:

R¹ and R² are independently hydrogen, halo, C₁-C₆ alkyl, hydroxy, or C₁-C₆ alkoxy;

15

R⁵, R⁶, and R⁷, are independently hydrogen, halo, C₁-C₆ alkyl, C₁-C₆ alkoxy, trifluoromethyl, or hydroxy;

R³ is hydrogen, C₂-C₇ alkanoyl, glycyl, or dimethylglycyl;

20

n is 1-6;

D is -S(O)_m-, -NH-, -C(O)-, or -O-,

25

m is 0, 1, or 2;

- 4 -

5 R⁸ and R⁹ are independently hydrogen, hydroxy, C₁-C₆ alkyl, C₁-C₆ alkoxy, C₁-C₆ alkylthio, C₁-C₆ alkoxy(C₁-C₆ alkylenyl)-, C₂-C₇ alkoxycarbonyl, C₂-C₇ alkoxy-
10 alkoxy(C₁-C₆ alkylenyl)-, trifluoromethoxy, trichloromethoxy, C₁-C₆ alkylthio, C₁-C₆ alkylamino, di(C₁-C₆ alkyl)amino, formyl, cyano, halo, trifluoromethyl, R¹⁰R¹¹N(C₁-C₆ alkylenyl)-, pyrrolyl, triazolyl, imidazolyl, tetrazolyl, thiazolyl, thiazolyl, thiadiazolyl, thiadiazolyl,
15 thiadiazolyl, piperidyl, pyrrolidyl, morpholinyl, morpholinocarbonyl, hexamethyleneiminyl, methylsulfonyl, methylsulfinyl, phenoxy, benzyloxy, carboxy, carbamoyl, or C₂-C₇ alkylcarbamoyl(C₁-C₆ alkylenyl)-,

15 R¹⁰ and R¹¹ are independently hydrogen or C₁-C₆ alkyl,

20 said C₁-C₆ alkyl or C₁-C₆ alkoxy groups being optionally substituted with one, two, or three moieties selected from the group consisting of hydroxy, halo, cyano, amino, nitro, carboxy, carbamoyl, and thiol;

25 or R⁸ and R⁹ can combine, together with the benzo ring to which they are attached, to form a naphthyl, dihydronaphthyl, tetrahydronaphthyl, quinolinyl, isoquinolinyl, 2-coumaranonyl, 3-coumaranonyl, benzothiazolyl, benzimidazolyl, indolyl, benzothienyl,
30 benzofuryl, 2,3-dihydrobenzofuryl, indolinyl, or 2,3-dihydrobenzothienyl group, which can be attached to D at any position of the bicyclic group;

or a pharmaceutically acceptable salt or solvate thereof.
35

- 5 -

In another embodiment this invention provides methods of treating a condition associated with an excess of tachykinins, which comprises administering to a mammal in need thereof an effective amount of a compound of Formula I, or a pharmaceutically acceptable salt or solvate thereof.

This invention also provides pharmaceutical formulations comprising, as an active ingredient, a compound of Formula I, or a pharmaceutically acceptable salt or solvate thereof, in combination with one or more pharmaceutically acceptable carriers, diluents, or excipients therefor.

Detailed Description and Preferred Embodiments

The terms and abbreviations used in the instant examples have their normal meanings unless otherwise designated. For example "°C" refers to degrees Celsius; "N" refers to normal or normality; "mol" refers to mole or moles; "mmol" refers to millimole or millimoles; "g" refers to gram or grams; "kg" refers to kilogram or kilograms; "L" refers to liter or liters; "ml" means milliliter or milliliters; "M" refers to molar or molarity; "MS" refers to mass spectrometry; and "NMR" refers to nuclear magnetic resonance spectroscopy.

"C₁-C₆ alkoxy" represents a straight or branched alkyl chain having from one to six carbon atoms attached to an oxygen atom. Typical C₁-C₆ alkoxy groups include methoxy, ethoxy, propoxy, isopropoxy, butoxy, *t*-butoxy, pentoxy and the like. The term "C₁-C₆ alkoxy" includes within its definition the terms "C₁-C₄ alkoxy" and "C₁-C₃ alkoxy".

As used herein, the term "C₁-C₁₂ alkyl" refers to straight or branched, monovalent, saturated aliphatic chains of 1 to 12 carbon atoms and includes, but is not limited to, methyl, ethyl, propyl, isopropyl, butyl, isobutyl, *t*-butyl, pentyl, isopentyl, and hexyl. The term "C₁-C₁₂ alkyl" includes within its definition the terms "C₁-C₆ alkyl" and "C₁-C₄ alkyl".

"C₂-C₇ alkanoyloxy" represents a straight or branched alkyl chain having from one to six carbon atoms attached to a carbonyl moiety joined through an oxygen atom. Typical C₂-C₇ alkanoyloxy groups

- 6 -

include acetoxy, propanoyloxy, isopropanoyloxy, butanoyloxy, *t*-butanoyloxy, pentanoyloxy, hexanoyloxy, 3-methylpentanoyloxy and the like.

5 "C₃-C₈ cycloalkyl" represents a saturated hydrocarbon ring structure containing from three to eight carbon atoms. Typical C₃-C₈ cycloalkyl groups include cyclopropyl, cyclopentyl, cyclohexyl, cycloheptyl, and the like.

"Halo" represents chloro, fluoro, bromo or iodo.

10 "C₁-C₆ alkylthio" represents a straight or branched alkyl chain having from one to six carbon atoms attached to a sulfur atom. Typical C₁-C₆ alkylthio groups include methylthio, ethylthio, propylthio, isopropylthio, butylthio and the like.

15 "C₁-C₁₂ alkylenyl" refers to a straight or branched, divalent, saturated aliphatic chain of 1 to 12 carbon atoms and includes, but is not limited to, methylenyl, ethylenyl, propylenyl, isopropylenyl, butylenyl, isobutylenyl, *t*-butylenyl, pentylenyl, isopentylenyl, hexylenyl, octylenyl, 3-methyloctylenyl, decylenyl. The term "C₁-C₆ alkylenyl" is encompassed within the term "C₁-C₁₂ alkylenyl".

20 "C₁-C₁₀ alkylamino" represents a group of the formula
-NH(C₁-C₁₀ alkyl)


wherein a chain having from one to ten carbon atoms is attached to an amino group. Typical C₁-C₄ alkylamino groups include methylamino, ethylamino, propylamino, isopropylamino, butylamino, *sec*-butylamino and the like.

25 "C₂-C₆ alkanoyl" represents a straight or branched alkyl chain having from one to five carbon atoms attached to a carbonyl moiety. Typical C₂-C₆ alkanoyl groups include ethanoyl (acetyl), propanoyl, isopropanoyl, butanoyl, *t*-butanoyl, pentanoyl, hexanoyl, 3-methylpentanoyl and the like.

30 "C₂-C₇ alkoxycarbonyl" represents a straight or branched alkoxy chain having from one to six carbon atoms attached to a carbonyl moiety. Typical C₂-C₇ alkoxycarbonyl groups include methoxycarbonyl, ethoxycarbonyl, propoxycarbonyl, isopropoxycarbonyl, butoxycarbonyl, *t*-butoxycarbonyl and the like.

- 7 -


The term "carbamoyl" as used herein refers to a moiety having one of the following structures.



5


The term "C₂-C₇ alkylcarbamoyl" as used herein refers to a branched or unbranched chain of 1 to 6 carbon atoms combined with a carbamoyl group, as such is defined above. This moiety has the following structure.

10



The term "haloformate" as used herein refers to an ester of a haloformic acid, this compound having the formula

15



wherein X is halo, and R^d is C₁-C₆ alkyl. Preferred haloformates are bromoformates and chloroformates. Especially preferred are chloroformates. Those haloformates wherein R^d is C₃-C₆ alkyl are especially preferred. Most preferred is isobutylchloroformate.

The compounds prepared in the processes of the present invention have an asymmetric center. As a consequence of this chiral center, the compounds produced in the present invention may occur as racemates, mixtures of enantiomers and as individual enantiomers, as well as diastereomers and mixtures of diastereomers.

The terms "R" and "S" are used herein as commonly used in organic chemistry to denote specific configuration of a chiral center. The term "R" (*rectus*) refers to that configuration of a chiral center with a clockwise relationship of group priorities (highest to second lowest) when viewed along the bond toward the lowest priority group. The term "S"

- 8 -

(*sinister*) refers to that configuration of a chiral center with a counterclockwise relationship of group priorities (highest to second lowest) when viewed along the bond toward the lowest priority group. The priority of groups is based upon their atomic number (in order of decreasing atomic number). A partial list of priorities and a discussion of stereochemistry is contained in NOMENCLATURE OF ORGANIC COMPOUNDS: PRINCIPLES AND PRACTICE, (J.H. Fletcher, *et al.*, eds., 1974) at pages 103-120.

In addition to the (R)-(S) system, the older D-L system is also used in this document to denote absolute configuration, especially with reference to amino acids. In this system a Fischer projection formula is oriented so that the number 1 carbon of the main chain is at the top. The prefix "D" is used to represent the absolute configuration of the isomer in which the functional (determining) group is on the right side of the carbon atom at the chiral center and "L", that of the isomer in which it is on the left.

The term "amino-protecting group" as used in the specification refers to substituents of the amino group commonly employed to block or protect the amino functionality while reacting other functional groups on the compound. Examples of such amino-protecting groups include formyl, trityl (herein abbreviated as "Tr"), phthalimido, trichloroacetyl, chloroacetyl, bromoacetyl, iodoacetyl, and urethane-type blocking groups such as benzyloxycarbonyl, 4-phenylbenzyloxycarbonyl, 2-methylbenzyloxycarbonyl, 4-methoxybenzyloxycarbonyl, 4-fluorobenzyloxycarbonyl, 4-chlorobenzyloxycarbonyl, 3-chlorobenzyloxycarbonyl, 2-chlorobenzyloxycarbonyl, 2,4-dichlorobenzyloxycarbonyl, 4-bromobenzyloxycarbonyl, 3-bromobenzyloxycarbonyl, 4-nitrobenzyloxycarbonyl, 4-cyanobenzyloxycarbonyl, *t*-butoxycarbonyl (herein abbreviated as "BoC"), 1,1-diphenyleth-1-yloxycarbonyl, 1,1-diphenylprop-1-yloxycarbonyl, 2-phenylprop-2-yloxycarbonyl, 2-(*p*-toluyl)-prop-2-yloxycarbonyl, cyclopentanyloxycarbonyl, 1-methylcyclopentanyloxycarbonyl, cyclohexanyloxycarbonyl, 1-methylcyclohexanyloxycarbonyl, 2-methylcyclohexanyloxycarbonyl, 2-(4-toluylsulfonyl)-ethoxycarbonyl, 2-(methylsulfonyl)ethoxycarbonyl,

- 9 -

2-(triphenylphosphino)-ethoxycarbonyl, fluorenylmethoxy-carbonyl ("Fmoc"), 2-(trimethylsilyl)ethoxycarbonyl, allyloxycarbonyl, 1-(trimethylsilylmethyl)prop-1-enyloxycarbonyl, 5-benzisoxalylmethoxycarbonyl, 4-acetoxycarboxybenzyloxycarbonyl, 2,2,2-trichloroethoxycarbonyl, 2-ethynyl-2-propoxycarbonyl, cyclopropylmethoxycarbonyl, 4-(decyloxy)benzyloxycarbonyl, isobornyloxycarbonyl, 1-piperidyloxycarbonyl and the like; benzoylmethylsulfonyl group, 2-nitrophenylsulfonyl, diphenylphosphine oxide and like amino-protecting groups. The species of amino-protecting group employed is usually not critical so long as the derivatized amino group is stable to the condition of subsequent reactions on other positions of the intermediate molecule and can be selectively removed at the appropriate point without disrupting the remainder of the molecule including any other amino-protecting groups. Preferred amino-protecting groups are trityl, *t*-butoxycarbonyl (*t*-BOC), allyloxycarbonyl and benzyloxycarbonyl. Further examples of groups referred to by the above terms are described by E. Haslam, "Protective Groups in Organic Chemistry", (J.G.W. McOmie, ed., 1973), at Chapter 2; and T.W. Greene and P.G.M. Wuts, PROTECTIVE GROUPS IN ORGANIC SYNTHESIS, (1991), at Chapter 7.

The term "carboxy-protecting group" as used in the specification refers to substituents of the carboxy group commonly employed to block or protect the carboxy functionality while reacting other functional groups on the compound. Examples of such carboxy-protecting groups include methyl, *p*-nitrobenzyl, *p*-methylbenzyl, *p*-methoxy-benzyl, 3,4-dimethoxybenzyl, 2,4-dimethoxybenzyl, 2,4,6-trimethoxybenzyl, 2,4,6-trimethylbenzyl, pentamethylbenzyl, 3,4-methylenedioxybenzyl, benzhydryl, 4,4'-dimethoxybenzhydryl, 2,2',4,4'-tetramethoxybenzhydryl, *t*-butyl, *t*-amyl, trityl, 4-methoxytrityl, 4,4'-dimethoxytrityl, 4,4',4''-trimethoxytrityl, 2-phenylprop-2-yl, trimethylsilyl, *t*-butyldimethylsilyl, phenacyl, 2,2,2-trichloroethyl, 2-(di(*n*-butyl)methylsilyl)ethyl, *p*-toluenesulfonyl, 4-nitrobenzylsulfonyl, allyl, cinnamyl, 1-(trimethylsilylmethyl)prop-1-en-3-yl and like moieties. Preferred carboxy-protecting groups are allyl, benzyl and *t*-butyl. Further examples

- 10 -

of these groups are found in E. Haslam, supra, at Chapter 5, and T.W. Greene, et al., supra, at Chapter 5.

The term "hydroxy-protecting groups" as used herein refers to substituents of the hydroxy group commonly employed to block or protect the hydroxy functionality while reacting other functional groups on the compound. Examples of such hydroxy-protecting groups include methoxymethyl, benzyloxymethyl, methoxyethoxymethyl, 2-(trimethylsilyl)ethoxymethyl, methylthiomethyl, 2,2-dichloro-1,1-difluoroethyl, tetrahydropyranyl, phenacyl, cyclopropylmethyl, allyl, C₁-C₆ alkyl, 2,6-dimethylbenzyl, o-nitrobenzyl, 4-picolyl, dimethylsilyl, *t*-butyldimethylsilyl, levulinate, pivaloate, benzoate, dimethylsulfonate, dimethylphosphinyl, isobutyrate, adamantate and tetrahydropyranyl. Further examples of these groups may be found in T. W. Greene and P.G.M. Wuts, PROTECTIVE GROUPS IN ORGANIC SYNTHESIS, (1991) at Chapter 3.

The term "leaving group" as used herein refers to a group of atoms that is displaced from a carbon atom by the attack of a nucleophile in a nucleophilic substitution reaction. The term "leaving group" as used in this document encompasses, but is not limited to, activating groups.

The term "activating group" as used herein refers a leaving group which, when taken with the carbonyl (-C=O) group to which it is attached, is more likely to take part in an acylation reaction than would be the case if the group were not present, as in the free acid. Such activating groups are well-known to those skilled in the art and may be, for example, succinimidoxo, phthalimidoxo, benzotriazolyloxy, benzenesulfonyloxy, methanesulfonyloxy, toluenesulfonyloxy, azido, or -O-CO-(C₄-C₇ alkyl).

As noted supra, this invention includes the pharmaceutically acceptable salts of the compounds defined by Formula I. A compound of this invention can possess a sufficiently acidic, a sufficiently basic, or both functional groups, and accordingly react with any of a number of organic and inorganic bases, and inorganic and organic acids, to form a pharmaceutically acceptable salt.

The term "pharmaceutically acceptable salt" as used herein, refers to salts of the compounds of the above formula which are

- 11 -

substantially non-toxic to living organisms. Typical pharmaceutically acceptable salts include those salts prepared by reaction of the compounds of the present invention with a pharmaceutically acceptable mineral or organic acid or an organic or inorganic base. Such salts are known as acid addition and base addition salts.

Acids commonly employed to form acid addition salts are inorganic acids such as hydrochloric acid, hydrobromic acid, hydroiodic acid, sulfuric acid, phosphoric acid, and the like, and organic acids such as *p*-toluenesulfonic acid, methanesulfonic acid, oxalic acid, *p*-bromophenylsulfonic acid, carbonic acid, succinic acid, citric acid, benzoic acid, acetic acid, and the like. Examples of such pharmaceutically acceptable salts are the sulfate, pyrosulfate, bisulfate, sulfite, bisulfite, phosphate, monohydrogenphosphate, dihydrogenphosphate, metaphosphate, pyrophosphate, bromide, iodide, acetate, propionate, decanoate, caprylate, acrylate, formate, hydrochloride, dihydrochloride, isobutyrate, caproate, heptanoate, propiolate, oxalate, malonate, succinate, suberate, sebacate, fumarate, maleate, butyne-1,4-dioate, hexyne-1,6-dioate, benzoate, chlorobenzoate, methylbenzoate, hydroxybenzoate, methoxybenzoate, phthalate, xylenesulfonate, phenylacetate, phenylpropionate, phenylbutyrate, citrate, lactate, γ -hydroxybutyrate, glycolate, tartrate, methanesulfonate, propanesulfonate, naphthalene-1-sulfonate, naphthalene-2-sulfonate, mandelate and the like. Preferred pharmaceutically acceptable acid addition salts are those formed with mineral acids such as hydrochloric acid and hydrobromic acid, and those formed with organic acids such as maleic acid and methanesulfonic acid.

Salts of amine groups may also comprise quaternary ammonium salts in which the amino nitrogen carries a suitable organic group such as an alkyl, alkenyl, alkynyl, or aralkyl moiety.

Base addition salts include those derived from inorganic bases, such as ammonium or alkali or alkaline earth metal hydroxides, carbonates, bicarbonates, and the like. Such bases useful in preparing the salts of this invention thus include sodium hydroxide, potassium hydroxide, ammonium hydroxide, potassium carbonate, sodium carbonate, sodium bicarbonate, potassium bicarbonate, calcium

- 12 -

hydroxide, calcium carbonate, and the like. The potassium and sodium salt forms are particularly preferred.

It should be recognized that the particular counterion forming a part of any salt of this invention is usually not of a critical nature, so long as the salt as a whole is pharmacologically acceptable and as long as the counterion does not contribute undesired qualities to the salt as a whole.

This invention further encompasses the pharmaceutically acceptable solvates of the compounds of Formulas I. Many of the Formula I compounds can combine with solvents such as water, methanol, ethanol and acetonitrile to form pharmaceutically acceptable solvates such as the corresponding hydrate, methanolate, ethanolate and acetonitrilate.


This invention also encompasses the pharmaceutically acceptable prodrugs of the compounds of Formula I. A prodrug is a drug which has been chemically modified and may be biologically inactive at its site of action, but which may be degraded or modified by one or more enzymatic or other *in vivo* processes to the parent bioactive form. This prodrug should have a different pharmacokinetic profile than the parent, enabling easier absorption across the mucosal epithelium, better salt formation or solubility, or improved systemic stability (an increase in plasma half-life, for example).

Typically, such chemical modifications include:

- 1) ester or amide derivatives which may be cleaved by esterases or lipases;
- 2) peptides which may be recognized by specific or nonspecific proteases; or
- 3) derivatives that accumulate at a site of action through membrane selection of a prodrug form or a modified prodrug form; or any combination of 1 to 3, *supra*. Conventional procedures for the selection and preparation of suitable prodrug derivatives are described, for example, in H, Bundgaard, DESIGN OF PRODRUGS, (1985).

The compounds of Formula I are generally prepared by reacting a compound of Formula II


- 13 -



II

- 5 with an appropriately substituted carboxylic acid, anhydride, or carboxylic acid halide in the presence of typical peptide coupling reagents such as N,N'-carbonyldiimidazole (CDI), N,N'-dicyclohexylcarbodiimide (DCC) and 1-(3-dimethylaminopropyl)-3-ethylcarbodiimide hydrochloride (EDC). A polymer supported form of EDC has been described
- 10 (Tetrahedron Letters, 34(48):7685 (1993)) and is very useful for the preparation of the compounds of the present invention. Isolation of products from reactions where a polymer bound reagent has been used is greatly simplified, requiring only filtration of the reaction mixture and then concentration of the filtrate under reduced pressure. The product
- 15 from these reactions may be purified chromatographically or recrystallized from a suitable solvent if desired..

Another preferred method of preparing the compounds of Formula I where D is -O- is by reacting a compound of Formula III




III

- 14 -

where X is a leaving group, preferably a halo moiety, most preferably a bromo group, with an appropriately substituted phenol, naphthol, or the like.


5 The most preferred method, to date, of synthesizing the
intermediates of Formulae II and III is depicted in Scheme I, infra. Many
of the steps of this synthesis are described in Patent Cooperation Treaty
Publication WO 95/14017, published May 26, 1995; European Patent
Application Publication 693,489, published January 24, 1996; and United
States Patent 5,530,009, issued June 25, 1996, the entire contents of
10 which are herein incorporated by reference.

- 15 -
Scheme I



wherein "Tr" refers to a trityl group, and "NMM" refers to N-methylmorpholine.


- 16 -

Scheme I (continued)

- 17 -


In another method of preparing the intermediates of Formulae II and III, Steps a) and b) can be combined as taught in United States Patent Application 60/021,849, filed July 16, 1996. In this method a compound of the formula

5



is prepared by reacting a compound of the formula

10



15

with bis(trimethylsilyl)amine in acetonitrile, followed by the addition of trityl chloride, N-methylmorpholine and 2-chloro-4,6-dimethoxy-1,3,5-triazine, in the presence of acetonitrile, and then adding 2-methoxybenzylamine.

The factor eventually found critical to the combination of the steps was the desilylation of the compound of Formula A, in Step (a), prior to ester formation via the addition of 2-chloro-4,6-dimethoxy-1,3,5-triazine (CDMT). In Step (a), desilylation had been accomplished by the addition of excess water prior to isolation which also dissolved any salts present.

20

- 18 -

When desilylating Formula A in the combined chemistry, stoichiometry becomes increasingly important. Consideration must be given to the presence of excess HMDS used in the initial silylation of D-tryptophan. Simply adding a stoichiometric amount of methyl alcohol (or water) relative to D-tryptophan will not allow the subsequent esterification to progress. Methyl alcohol must also be added to quench all remaining unreacted HMDS. However, any excess methyl alcohol will consume CDMT and prevent complete esterification. Once the desilylation of the compound of Formula A and decomposition of excess HMDS is complete, the chemistry of Step (b) proceeds as expected and high quality desired intermediate is produced in good yield.

In the above process, the intermediate amides are reduced to amines using procedures well known in the art. These reductions can be performed using lithium aluminum hydride as well as by use of many other different aluminum-based hydrides. An especially preferred reagent employed in this reduction is RED-AL[®], which is the tradename of a 3.4 M solution of sodium bis(2-methoxyethoxy)aluminum hydride in toluene. Alternatively, the amides can be reduced by catalytic hydrogenation, though high temperatures and pressures are usually required for this. Sodium borohydride in combination with other reagents may be used to reduce the amide. Borane complexes, such as a borane dimethylsulfide complex, are especially useful in this reduction reaction.

The acylation of the secondary amine can be done using any of a large number of techniques regularly employed by those skilled in organic chemistry. One such reaction scheme is a substitution using an anhydride such as acetic anhydride. Another reaction scheme often employed to acylate a secondary amine employs a carboxylic acid preferably with an activating agent. An amino-de-alkoxylation type of reaction uses esters as a means of acylating the amine. Activated esters which are attenuated to provide enhanced selectivity are very efficient acylating agents. One preferred such activated ester is *p*-nitrophenyl ester, such as *p*-nitrophenyl acetate.

- 19 -

Primary amines can also be acylated using amides to perform what is essentially an exchange reaction. This reaction is usually carried out with the salt of the amine. Boron trifluoride, usually in the form of a boron trifluoride diethyl ether complex, is frequently added to this reaction to complex with the leaving ammonia.

An additional step is one of substitution of the secondary amine. For most of the compounds of Formula I this substitution is one of alkylation, acylation, or sulfonation. This substitution is usually accomplished using well recognized means. Typically, alkylations can be achieved using alkyl halides and the like as well as the well-known reductive alkylation methods, employing aldehydes or ketones. Many of the acylating reaction protocols discussed *supra* efficiently acylate the secondary amine as well. Alkyl- and aryl-sulfonyl chlorides can be employed to sulfonate the secondary amine.

In many instances one of the later steps in the synthesis of the compounds of Formulae II and III is the removal of an amino- or carboxy-protecting group. Such procedures, which vary, depending upon the type of protecting group employed as well as the relative lability of other moieties on the compound, are described in detail in many standard references works such as T.W. Greene, *et al.*, PROTECTIVE GROUPS IN ORGANIC SYNTHESIS (1991).

The following Examples and Preparations further illustrate the compounds of the present invention and the methods for their synthesis. The Examples are not intended to be limiting to the scope of the invention in any respect, and should not be so construed. All experiments were run under a positive pressure of dry nitrogen or argon. All solvents and reagents were purchased from commercial sources and used as received, unless otherwise indicated. Dry tetrahydrofuran (THF) was obtained by distillation from sodium or sodium benzophenone ketyl prior to use.

Proton nuclear magnetic resonance (^1H NMR) spectra were obtained on a GE QE-300 spectrometer at 300.15 MHz, a Bruker AM-500 spectrometer at 500 MHz, or a Bruker AC-200P spectrometer at 200 MHz.

- 20 -

(Unless designated otherwise, the term "NMR" as employed herein refers to proton nuclear magnetic resonance.) Free atom bombardment mass spectroscopy (FAB) was performed on a VG ZAB-2SE instrument. Field desorption mass spectroscopy (FDMS) was performed using either a VG
5 70SE or a Varian MAT 731 instrument.

Optical rotations were measured with a Perkin-Elmer 241 polarimeter. Chromatographic separation on a Waters Prep 500 LC was generally carried out using a linear gradient of the solvents indicated in the text unless otherwise specified.

10 The reactions were generally monitored for completion using thin layer chromatography (TLC). Thin layer chromatography was performed using E. Merck Kieselgel 60 F₂₅₄ plates, 5 cm x 10 cm, 0.25 mm thickness. Spots were detected using a combination of UV and chemical detection (plates dipped in a ceric ammonium molybdate solution [75 g of
15 ammonium molybdate and 4 g of cerium (IV) sulfate in 500 ml of 10% aqueous sulfuric acid] and then heated on a hot plate). Preparative centrifugal thin layer chromatography was performed on a Harrison Model 7924A Chromatotron using Analtech silica gel GF rotors.

Cation exchange chromatography was performed with
20 Dowex® 50X8-100 ion exchange resin. Anion exchange chromatography was performed with Bio-Rad AG® 1-X8 anion-exchange resin (acetate form converted to hydroxide form). Flash chromatography was performed as described by Still, *et al.*, Journal of Organic Chemistry, 43:2923 (1978).

Optical rotations are reported at the sodium-D-line (354 nm).
25 Elemental analyses for carbon, hydrogen, and nitrogen were determined on a Control Equipment Corporation 440 Elemental Analyzer, or were performed by the Universidad Complutense Analytical Centre (Facultad de Farmacia, Madrid, Spain). Melting points were determined in open glass capillaries on a Thomas Hoover capillary melting point apparatus or
30 a Büchi melting point apparatus, and are uncorrected.

The following methods provide illustrative protocols for preparing the compounds of Formula I as depicted in the Schemes supra. Throughout the Methods and Examples, *infra*, the terms "NMR", "IR",


- 21 -

and "UV" indicate that the proton nuclear magnetic resonance, infrared, and ultraviolet spectroscopy, respectively, were consistent with the desired title product.

5

Preparation 1

Preparation of (R)-3-(1H-indol-3-yl)-N-(2-methoxybenzyl)-2-(N-triphenylmethylamino)propanamide.



10

In a 50 gallon, glass-lined reactor, L-tryptophan (4.50 kg, 22.0 mol) was added to acetonitrile (30 L, 6.7 vol) at 20 °C. This reactor was vented to a scrubber containing water, intended to scrub ammonia generated during the silylation reaction and HCl generated during the tritylation and esterification reactions. Bis(trimethylsilyl)amine (HMDS, 5.81 L, 27.5 mol, 1.25 eq) was transferred by gravity to the L-tryptophan slurry from a plastic carboy. The carboy was rinsed with acetonitrile (0.5 L). The slurry was heated to 55 °C and stirred until reaction completion. The reaction endpoint was defined as the point at which the slurry has completely gone into solution. The reaction was clear yellow at completion and took about 2 hours.

Trityl chloride (6.45 kg, 23.1 mol, 1.05 eq) was slurried in acetonitrile (30 L, 6.7 vol) and transferred into the reactor at 47 °C, using trapped vacuum at 325 mm Hg.

25

- 22 -

N-methylmorpholine (5.38 L, 48.9 mol, 2.20 eq) was also transferred into the reactor at this time. The reaction slurry was heated and maintained at 55 °C until reaction completion, determined by high performance liquid chromatography analysis. Reaction time was about 2.5 hours.

The reactor was isolated from the scrubber, and cooled to 35-40 °C. Methyl alcohol (2.29 L, 56.5 mol, 2.55 eq.) was charged to the reactor and the mixture cooled to 25 °C. 2-Chloro-4,6-dimethoxy-1,3,5-triazine (CDMT, 4.14 kg, 23.61 mol, 1.07 eq) was added to the reactor with acetonitrile (28 L, 6.2 vol) at 25 °C. The reactor was again vented to the scrubber. The reaction slurry was stirred at room temperature until completion. The reaction endpoint is determined by high performance liquid chromatography analysis. Reaction time is approximately 2 hours. The reactor was isolated from the scrubber following the reaction.

2-Methoxybenzylamine (3.11 L, 23.8 mol, 1.08 eq) was charged to the reactor from a plastic carboy by gravity. The slurry thickens with the addition of 2-methoxybenzylamine. The reaction slurry was heated to 35°C and stirred until reaction completion, determined by high performance liquid chromatography analysis. Reaction time was 2.5 hours.

Water (45 kg, 10 vol) was pre-weighed into a separate 50 gallon, glass-lined tank. The water was pressure-transferred into the reaction mixture slurry over about 45 minutes. The resulting yellow-colored slurry was cooled to 0-5 °C over two hours and stirred overnight.

The title intermediate was isolated by vertical basket centrifuge isolation using three micron polyethylene multiple filament isolation bag. During the centrifugation, the load speed was generally between 900-1050 rpm, the wash speed was 900-1500 rpm, and the spin speed was 1500-2300 rpm.


The title intermediate was then dried by rotary vacuum drying. Yield: 86.4% with isomer purity of 99.6%.

Preparation 2

- 23 -

Reduction of Carbonyl

Preparation of (R)-3-(1H-indol-3-yl)-1-[N-(2-methoxybenzyl)amino]-2-(N-triphenylmethylamino)propane



RED-AL[®], [a 3.4 M, solution of sodium bis(2-methoxyethoxy)aluminum hydride in toluene] (535 ml, 1.819 mol), dissolved in anhydrous tetrahydrofuran (400 ml) was slowly added using an addition funnel to a refluxing solution of the acylation product, (R)-3-(1H-indol-3-yl)-N-(2-methoxybenzyl)-2-(N-triphenylmethylamino)propanamide (228.6 g, 0.404 mol) produced *supra*, in anhydrous tetrahydrofuran (1.0 L) under a nitrogen atmosphere. The reaction mixture became a purple solution. The reaction was quenched after at least 20 hours by the slow addition of excess saturated Rochelle's salt solution (potassium sodium tartrate tetrahydrate). The organic layer was isolated, washed with brine (2X), dried over anhydrous sodium sulfate, filtered, and concentrated to an oil on a rotary evaporator. No further purification was done and the product was used directly in the next step.


Preparation 3

- 24 -

Acylation of Secondary Amine

Preparation of (R)-3-(1H-indol-3-yl)-1-[N-(2-methoxybenzyl)-acetylaminol]-2-(N-triphenylmethylamino)propane

5



To a stirring solution of (R)-3-(1H-indol-3-yl)-1-[N-(2-methoxybenzyl)amino]-2-(N-triphenylmethylamino)propane (0.404 mol) in anhydrous tetrahydrofuran (1.2 L) under a nitrogen atmosphere at 0°C was added triethylamine (66.5 ml, 0.477 mol) and acetic anhydride (45.0 ml, 0.477 mol). After 4 hours, the mixture was concentrated on a rotary evaporator, redissolved in methylene chloride and ethyl acetate, washed with water (2X) and brine (2X), dried over anhydrous sodium sulfate, filtered, and concentrated to a solid on a rotary evaporator. The resulting solid was dissolved in chloroform and loaded onto silica gel 60 (230-400 mesh) and eluted with a 1:1 mixture of ethyl acetate and hexanes. The product was then crystallized from an ethyl acetate/hexanes mixture. The resulting product of (R)-3-(1H-indol-3-yl)-1-[N-(2-methoxybenzyl)acetylaminol]-2-(N-triphenylmethylamino)propane was crystallized and isolated over three crops giving 208.97 grams (87% yield) of analytically pure material.

Analysis for $C_{40}H_{39}N_3O_2$:

Theory: C, 80.91; H, 6.62; N, 7.08.


Found: C, 81.00; H, 6.69; N, 6.94.

- 25 -

Preparation 4

Deprotection

5



Preparation of (R)-2-amino-3-(1H-indol-3-yl)-1-[N-(2-methoxybenzyl)acetyl]amino]propane dihydrochloride

10

A stirring solution of (R)-3-(1H-indol-3-yl)-1-[N-(2-methoxybenzyl)acetyl]amino]-2-(N-triphenylmethylamino)propane in two volumes of methylene chloride was cooled to between -40°C and -50°C. Anhydrous hydrogen chloride gas was added at such a rate that the temperature of the reaction mixture did not exceed 0°C. The reaction mixture was stirred for 30 minutes to one hour at 0-10°C.

15

To this reaction mixture was added two volumes of methyl *t*-butyl ether and the resulting mixture was allowed to stir for 30 minutes to one hour at 0-10°C. The resulting crystalline solid was removed by filtration and then washed with methyl *t*-butyl ether. The reaction product was dried under vacuum at 50°C. (Yield >98%)

20

Analysis for $\text{C}_{21}\text{H}_{25}\text{N}_3\text{O}_2 \cdot 2 \text{ HCl}$:

Theory: C, 59.44; H, 6.41; N, 9.90.


Found: C, 60.40; H, 6.60; N, 9.99.

25

Preparation 5

Preparation of (R)-2-[(2-bromo)acetyl]amino-3-(1H-indol-3-yl)-1-[N-(2-methoxybenzyl)acetyl]amino]propane

- 26 -



To a stirring solution of (R)-2-amino-3-(1H-indol-3-yl)-1-[N-(2-methoxybenzyl)acetyl]propane (7.51 g, 21.369 mmol) in anhydrous tetrahydrofuran (100 ml) under a nitrogen atmosphere at 0°C were added diisopropylethylamine (4.1 ml, 23.537 mmol) and bromoacetyl bromide (2.05 ml, 23.530 mmol). After 2 hours, ethyl acetate was added and the reaction mixture washed with water twice, 1.0 N hydrochloric acid (2X), saturated sodium bicarbonate solution (2X), and brine. The organic layer was dried over anhydrous sodium sulfate, filtered, and concentrated to a tan foam on a rotary evaporator. In this manner the 2-[(2-bromo)acetyl]amino-3-(1H-indol-3-yl)-1-[N-(2-methoxybenzyl)acetyl]propane was obtained in quantitative yield. No further purification was necessary.

Preparation 6

Preparation of a polystyrene bound isocyanate resin

20

To a stirred suspension of 50 grams (61 mmol) aminomethylated polystyrene resin (1.22 mmol/g) in 800 ml toluene was added 193 ml (366 mmol) 1.9 M phosgene in toluene. After stirring the reaction mixture for 10 minutes, 67 ml (482 mmol) triethylamine was added and the reaction mixture was stirred for 18 hours at room temperature. The mixture was filtered and the recovered solid washed with 10 times with dichloromethane. A light pink resin mixed with a white solid was obtained. This solid mixture

- 27 -

was resuspended in 700 ml dichloromethane, stirred for 10 minutes and then filtered and washed well with dichloromethane. The resulting solid was again suspended, stirred and washed with dichloromethane to provide the desired resin.

5 IR(KBr): 2252 cm^{-1} (characteristic peak for -N=C=O)

General Procedure I

To a suspension of three equivalents of polymer-bound 1-piperidine in 1 ml of chloroform (R)-2-amino-3-(1H-indol-3-yl)-1-[N-(2-methoxybenzyl)acetylaminopropyl]propane dihydrochloride (10 mg, 0.024 mmol, 1 eq) was dissolved. To this mixture was added the appropriate carboxylic acid (0.036 mmol, 1.5 eq) and polymer-bound 1-(3-dimethylaminopropyl)-3-propylcarbodiimide hydrochloride (108 mg, 0.108 mmol, 4.5 eq). The resulting mixture was agitated at ambient temperature for about 2-3 days. Unreacted (R)-2-amino-3-(1H-indol-3-yl)-1-[N-(2-methoxybenzyl)acetylaminopropyl]propane dihydrochloride was removed by the addition of an excess of polystyrene bound isocyanate resin and rocked for four hours. The reaction mixtures were filtered and the filtrates were concentrated.

General Procedure II


In a reaction vial potassium tert-butoxide (19.38 mg, 0.158 mmol, 3 eq) and an appropriate substituted phenol or naphthol (0.158 mmol, 3 eq) were admixed in 0.7 ml of dry tetrahydrofuran. To this mixture (R)-2-[(2-bromo)acetyl]amino-3-(1H-indol-3-yl)-1-[N-(2-methoxybenzyl)acetylaminopropyl]propane (25 mg, 0.053 mmol, 1 eq) was added and the resulting mixture was heated for two hours at 80°C. The solvents were removed in vacuo and the residue was redissolved in methylene chloride and washed once with water. The organic fraction was dried over sodium sulfate. The solvents were removed in vacuo.

- 28 -

Example 1

Preparation of (R)-2-[(2-phenoxy)acetyl]amino-3-(1H-indol-3-yl)-1-[N-(2-methoxybenzyl)acetyl]amino]propane

5




NMR was consistent with the proposed title structure.

10

Example 2

Preparation of (R)-2-[[2-(4-methoxymethyl)phenoxy]acetyl]amino-3-(1H-indol-3-yl)-1-[N-(2-methoxybenzyl)acetyl]amino]propane

- 29 -




NMR was consistent with the proposed title structure.

5

Example 3

Preparation of (R)-2-[[2-(4-carboxy)phenoxy]acetyl]amino-3-(1H-indol-3-yl)-1-[N-(2-methoxybenzyl)acetyl]amino]propane




10

NMR was consistent with the proposed title structure.

- 30 -

Example 4

Preparation of (R)-2-[[2-(4-ethoxy)phenoxy]acetyl]amino-3-(1H-indol-3-yl)-
5 1-[N-(2-methoxybenzyl)acetyl]amino]propane



NMR was consistent with the proposed title structure.


10

Example 5

Preparation of (R)-2-[[2-(4-methoxycarbonyl)phenoxy]acetyl]amino-3-(1H-
indol-3-yl)-1-[N-(2-methoxybenzyl)acetyl]amino]propane

15

- 31 -




NMR was consistent with the proposed title structure.

5

Example 6

Preparation of (R)-2-[[2-(4-trifluoromethoxy)phenoxy]acetyl]amino-3-(1H-indol-3-yl)-1-[N-(2-methoxybenzyl)acetyl]propane




10

- 32 -

NMR was consistent with the proposed title structure.

Example 7


- 5 Preparation of (R)-2-[[2-[4-[[[(methoxy)carbonyl]methyl]]phenoxy]acetyl]-amino-3-(1H-indol-3-yl)-1-[N-(2-methoxybenzyl)acetyl]amino]propane



- 10 NMR was consistent with the proposed title structure.

Example 8

- Preparation of (R)-2-[[2-[4-[[[(methoxy)carbonyl]ethyl]]phenoxy]acetyl]-
15 amino-3-(1H-indol-3-yl)-1-[N-(2-methoxybenzyl)acetyl]amino]propane




NMR was consistent with the proposed title structure.

5

Example 9

Preparation of (R)-2-[2-(naphth-2-yloxy)acetyl]amino-3-(1H-indol-3-yl)-1-[N-(2-methoxybenzyl)acetamino]propane



10


- 34 -

NMR was consistent with the proposed title structure.

Example 10

5

Preparation of (R)-2-[2-(4-acetylphenoxy)acetyl]-amino-3-(1H-indol-3-yl)-1-[N-(2-methoxybenzyl)acetyl]amino]propane



10


NMR was consistent with the proposed title structure.

Example 11

15

Preparation of (R)-2-[2-(4-hydroxyphenoxy)acetyl]-amino-3-(1H-indol-3-yl)-1-[N-(2-methoxybenzyl)acetyl]amino]propane

- 35 -




NMR was consistent with the proposed title structure.

5

Example 12

Preparation of (R)-2-[2-(4-chlorophenoxy)acetyl]-amino-3-(1H-indol-3-yl)-1-[N-(2-methoxybenzyl)acetyl]propane




10

NMR was consistent with the proposed title structure.

Example 13

- 36 -

Preparation of (R)-2-[2-[4-(benzyloxy)phenoxy]acetyl]-amino-3-(1H-indol-3-yl)-1-[N-(2-methoxybenzyl)acetyl]amino]propane




5

NMR was consistent with the proposed title structure.

Example 14

10

Preparation of (R)-2-[2-(4-hydroxymethylphenoxy)acetyl]-amino-3-(1H-indol-3-yl)-1-[N-(2-methoxybenzyl)acetyl]amino]propane




NMR was consistent with the proposed title structure.

5

Example 15

Preparation of (R)-2-[2-(4-methoxyphenoxy)acetyl]-amino-3-(1H-indol-3-yl)-1-[N-(2-methoxybenzyl)acetyl]amino]propane




10

NMR was consistent with the proposed title structure.

Example 16

- 38 -

Preparation of (R)-2-[2-(3-methoxyphenoxy)acetyl]-amino-3-(1H-indol-3-yl)-1-[N-(2-methoxybenzyl)acetylamino]propane




5

NMR was consistent with the proposed title structure.

Example 17

10

Preparation of (R)-2-[2-(2-methoxyphenoxy)acetyl]-amino-3-(1H-indol-3-yl)-1-[N-(2-methoxybenzyl)acetylamino]propane



15


NMR was consistent with the proposed title structure.

- 39 -

Example 18

Preparation of (R)-2-[2-(2-chlorophenoxy)acetyl]-amino-3-(1H-indol-3-yl)-1-[N-(2-methoxybenzyl)acetylaminopropyl]propane

5




NMR was consistent with the proposed title structure.

10

Example 19

Preparation of (R)-2-[2-(2-carbamoylphenoxy)acetyl]-amino-3-(1H-indol-3-yl)-1-[N-(2-methoxybenzyl)acetylaminopropyl]propane




15

- 40 -

NMR was consistent with the proposed title structure.

Example 20

- 5 Preparation of (R)-2-[2-(3,4-dichlorophenoxy)acetyl]-amino-3-(1H-indol-3-yl)-1-[N-(2-methoxybenzyl)acetyl]amino]propane




- 10 NMR was consistent with the proposed title structure.

Example 21

- Preparation of (R)-2-[2-(4-fluorophenoxy)acetyl]-amino-3-(1H-indol-3-yl)-
15 1-[N-(2-methoxybenzyl)acetyl]amino]propane

- 41 -




NMR was consistent with the proposed title structure.

5

Example 22

Preparation of (R)-2-[2-(4-nitrophenoxy)acetyl]-amino-3-(1H-indol-3-yl)-1-[N-(2-methoxybenzyl)acetyl]amino]propane




10

NMR was consistent with the proposed title structure.

Example 23

- 42 -

Preparation of (R)-2-[2-[4-(2-hydroxyethyl)phenoxy]acetyl]-amino-3-(1H-indol-3-yl)-1-[N-(2-methoxybenzyl)acetyl]amino]propane




NMR was consistent with the proposed title structure.

Example 24

10

Preparation of 2-[2-[4-(3-hydroxypropyl)phenoxy]acetyl]-amino-3-(1H-indol-3-yl)-1-[N-(2-methoxybenzyl)acetyl]amino]propane




- 43 -

NMR was consistent with the proposed title structure.

Example 25

- 5 Preparation of 2-[2-[4-(2-carbamoylethyl)phenoxy]acetyl]-amino-3-(1H-indol-3-yl)-1-[N-(2-methoxybenzyl)acetylaminopropyl]propane




- 10 NMR was consistent with the proposed title structure.

Example 26

- Preparation of 2-[2-[4-(triazol-1-yl)phenoxy]acetyl]-amino-3-(1H-indol-3-yl)-1-[N-(2-methoxybenzyl)acetylaminopropyl]propane
- 15

- 44 -




NMR was consistent with the proposed title structure.

5

Example 27

Preparation of 2-[2-[4-(ethyl)phenoxy]acetyl]-amino-3-(1H-indol-3-yl)-1-[N-(2-methoxybenzyl)acetyl]amino]propane



10


NMR was consistent with the proposed title structure.

- 45 -

Example 28

Preparation of 2-[2-[4-(*n*-propyl)phenoxy]acetyl]-amino-3-(1H-indol-3-yl)-
1-[N-(2-methoxybenzyl)acetyl]amino]propane

5




NMR was consistent with the proposed title structure.

10

Example 29

Preparation of 2-[2-[2-(hydroxymethyl)phenoxy]acetyl]-amino-3-(1H-indol-3-yl)-
1-[N-(2-methoxybenzyl)acetyl]amino]propane




15

- 46 -

NMR was consistent with the proposed title structure.

Example 30

- 5 Preparation of 2-[2-[4-(2-methoxyethyl)phenoxy]acetyl]-amino-3-(1H-indol-3-yl)-1-[N-(2-methoxybenzyl)acetyl]amino]propane




- 10 NMR was consistent with the proposed title structure.

Example 31

- Preparation of 2-[2-[4-[(morpholin-4-yl)carbonyl]phenoxy]acetyl]-amino-3-(1H-indol-3-yl)-1-[N-(2-methoxybenzyl)acetyl]amino]propane
- 15

- 47 -




NMR was consistent with the proposed title structure.

5

Example 32

Preparation of 2-[2-[4-(4-hydroxybutyl)phenoxy]acetyl]-amino-3-(1H-indol-3-yl)-1-[N-(2-methoxybenzyl)acetylaminopropyl]propane




10

- 48 -

NMR was consistent with the proposed title structure.

Example 33

- 5 Preparation of 2-[2-[2-coumaranon-5-yl-oxy]acetyl]-amino-3-(1H-indol-3-yl)-1-[N-(2-methoxybenzyl)acetylamino]propane




- 10 NMR was consistent with the proposed title structure.

Example 34

- Preparation of 2-[2-[benzothiazol-6-yl-oxy]acetyl]-amino-3-(1H-indol-3-yl)-
15 1-[N-(2-methoxybenzyl)acetylamino]propane

- 49 -




NMR was consistent with the proposed title structure.

5

Example 35

Preparation of 2-[2-[4-(methylsulfonyl)phenoxy]acetyl]-amino-3-(1H-indol-3-yl)-1-[N-(2-methoxybenzyl)acetylamino]propane




10

NMR was consistent with the proposed title structure.

Example 35

- 50 -

Preparation of 2-[2-[4-(methylsulfonyl)phenoxy]acetyl]-amino-3-(1H-indol-3-yl)-1-[N-(2-methoxybenzyl)acetyl]amino]propane



5


NMR was consistent with the proposed title structure.

Example 36

10

Preparation of 2-[2-[quinolin-6-yl]acetyl]-amino-3-(1H-indol-3-yl)-1-[N-(2-methoxybenzyl)acetyl]amino]propane

- 51 -




NMR was consistent with the proposed title structure.

5

Example 37

Preparation of 2-[2-[4-(3-cyanopropyl)phenoxy]acetyl]-amino-3-(1H-indol-3-yl)-1-[N-(2-methoxybenzyl)acetyl]propane




10

NMR was consistent with the proposed title structure.

Example 38

- 52 -

Preparation of 2-[2-[4-(imidazol-1-yl)phenoxy]acetyl]-amino-3-(1H-indol-3-yl)-1-[N-(2-methoxybenzyl)acetyl]amino]propane




5

NMR was consistent with the proposed title structure.

Example 39

10

Preparation of 2-[2-[4-(cyanomethyl)phenoxy]acetyl]-amino-3-(1H-indol-3-yl)-1-[N-(2-methoxybenzyl)acetyl]amino]propane




15

- 53 -

NMR was consistent with the proposed title structure.

Example 40

- 5 Preparation of 2-[2-[4-(isopropyl)phenoxy]acetyl]-amino-3-(1H-indol-3-yl)-
1-[N-(2-methoxybenzyl)acetyl]amino]propane




- 10 NMR was consistent with the proposed title structure.

Example 41

- Preparation of 2-[2-[3-(hydroxymethyl)phenoxy]acetyl]-amino-3-(1H-
15 indol-3-yl)-1-[N-(2-methoxybenzyl)acetyl]amino]propane

- 54 -




NMR was consistent with the proposed title structure.

5

Example 42

Preparation of 2-[2-[4-[[[(methylamino)carbonyl]methyl]phenoxy]acetyl]-amino-3-(1H-indol-3-yl)]-1-[N-(2-methoxybenzyl)acetyl]amino]propane



10


NMR was consistent with the proposed title structure.

Example 43

15

- 55 -

Preparation of 2-[2-[4-(formyl)phenoxy]acetyl]-amino-3-(1H-indol-3-yl)-1-[N-(2-methoxybenzyl)acetyl]amino]propane




5

NMR was consistent with the proposed title structure.

Example 44

10 Preparation of 2-[2-[4-(thiazolin-2-yl)phenoxy]acetyl]-amino-3-(1H-indol-3-yl)-1-[N-(2-methoxybenzyl)acetyl]amino]propane




- 56 -

NMR was consistent with the proposed title structure.

Example 45

- 5 Preparation of 2-[2-[4-(cyano)phenoxy]acetyl]-amino-3-(1H-indol-3-yl)-1-[N-(2-methoxybenzyl)acetyl]amino]propane




- 10 NMR was consistent with the proposed title structure.

Example 46

- Preparation of 2-[2-[4-(3-methyl-2-pyrazolin-5-yl)phenoxy]acetyl]-amino-3-(1H-indol-3-yl)-1-[N-(2-methoxybenzyl)acetyl]amino]propane
- 15

- 57 -




NMR was consistent with the proposed title structure.

5

Example 47

Preparation of 2-[2-[4-[2-(acetyl)ethyl]phenoxy]acetyl]-amino-3-(1H-indol-3-yl)-1-[N-(2-methoxybenzyl)acetyl]amino]propane




10

- 58 -

NMR was consistent with the proposed title structure.

Example 48

- 5 Preparation of 2-[2-[3-(2-hydroxyethyl)phenoxy]acetyl]-amino-3-(1H-indol-3-yl)-1-[N-(2-methoxybenzyl)acetyl]amino]propane




- 10 NMR was consistent with the proposed title structure.

Example 49

- Preparation of 2-[2-[4-[[[butylamino)carbonyl]amino]phenoxy]acetyl]-
15 amino-3-(1H-indol-3-yl)-1-[N-(2-methoxybenzyl)acetyl]amino]propane

- 59 -




NMR was consistent with the proposed title structure.

5

Example 50

Preparation of 2-[2-[4-(3,5-dichlorophenoxy)phenoxy]acetyl]-amino-3-(1H-indol-3-yl)-1-[N-(2-methoxybenzyl)acetyl]amino]propane




10

- 60 -

NMR was consistent with the proposed title structure.

Example 51

- 5 Preparation of 2-[2-[4-(acetamido)phenoxy]acetyl]-amino-3-(1H-indol-3-yl)-1-[N-(2-methoxybenzyl)acetyl]amino]propane




- 10 NMR was consistent with the proposed title structure.

Example 52

- Preparation of 2-[2-[4-(acetoxo)phenoxy]acetyl]-amino-3-(1H-indol-3-yl)-1-[N-(2-methoxybenzyl)acetyl]amino]propane
- 15

- 61 -




NMR was consistent with the proposed title structure.

5

Example 53

Preparation of 2-[3-(phenylsulfonyl)propanoyl]amino-3-(1H-indol-3-yl)-1-[N-(2-methoxybenzyl)acetyl]amino]propane




10

NMR was consistent with the proposed title structure.

- 62 -

Example 54

Preparation of 2-[4-(2,4-dichlorophenoxy)butanoyl]amino-3-(1H-indol-3-yl)-1-[N-(2-methoxybenzyl)acetyl]amino]propane




NMR was consistent with the proposed title structure.

Example 55

Preparation of 2-[3-[benzothiazol-2-ylthio]propanoyl]-amino-3-(1H-indol-3-yl)-1-[N-(2-methoxybenzyl)acetyl]amino]propane

- 63 -




NMR was consistent with the proposed title structure.

5

Example 56

Preparation of 2-[2-[4-(imidazol-1-yl)phenoxy]acetyl]-amino-3-(1H-indol-3-yl)-1-[N-(2-methoxybenzyl)acetylamino]propane




- 64 -

NMR was consistent with the proposed title structure.

Example 57

5

Preparation of 2-[2-[4-(imidazol-1-yl)phenoxy]acetyl]-amino-3-(1H-indol-3-yl)-1-[N-(2-chlorobenzyl)acetyl]amino]propane



10


NMR was consistent with the proposed title structure.

Example 57

15

Preparation of 2-[2-[4-(1,2,3-thiadiazol-4-yl)phenoxy]acetyl]-amino-3-(1H-indol-3-yl)-1-[N-(2-methoxybenzyl)acetyl]amino]propane

- 65 -




NMR was consistent with the proposed title structure. MS-FD 569 (M+).

5

Example 58

Preparation of 2-[3-[benzoyl]ethanoyl]-amino-3-(1H-indol-3-yl)-1-[N-(2-methoxybenzyl)acetyl]propane

- 66 -



NMR was consistent with the proposed title structure. MS-FD 569 (M+).

5 The compounds of the present invention possess tachykinin
receptor activity. The biological efficacy of a compound believed to be
effective as a tachykinin receptor antagonist may be confirmed by
employing an initial screening assay which rapidly and accurately
measured the binding of the tested compound to known NK-1 and NK-2
10 receptor sites. Assays useful for evaluating tachykinin receptor
antagonists are well known in the art. See, e.g., J. Jukic, et al., Life
Sciences, 49:1463-1469 (1991); N. Kucharczyk, et al., Journal of Medicinal
Chemistry, 36:1654-1661 (1993); N. Rouissi, et al., Biochemical and
Biophysical Research Communications, 176:894-901 (1991).

15

NK-1 Receptor Binding Assay

 Radioreceptor binding assays were performed using a
derivative of a previously published protocol. D.G. Payan, et al., Journal
20 of Immunology, 133:3260-3265 (1984). In this assay an aliquot of IM9
cells (1×10^6 cells/tube in RPMI 1604 medium supplemented with 10%
fetal calf serum) was incubated with 20 pM ^{125}I -labeled substance P in

- 67 -

the presence of increasing competitor concentrations for 45 minutes at 4°C.

The IM9 cell line is a well-characterized cell line which is readily available to the public. See, e.g., Annals of the New York Academy of Science, 190: 221-234 (1972); Nature (London), 251:443-444 (1974); Proceedings of the National Academy of Sciences (USA), 71:84-88 (1974). These cells were routinely cultured in RPMI 1640 supplemented with 50 µg/ml gentamicin sulfate and 10% fetal calf serum.

The reaction was terminated by filtration through a glass fiber filter harvesting system using filters previously soaked for 20 minutes in 0.1% polyethylenimine. Specific binding of labeled substance P was determined in the presence of 20 nM unlabeled ligand.

Many of the compounds employed in the methods of the present invention are also effective antagonists of the NK-2 receptor.

NK-2 Receptor Binding Assay

The CHO-hNK-2R cells, a CHO-derived cell line transformed with the human NK-2 receptor, expressing about 400,000 such receptors per cell, were grown in 75 cm² flasks or roller bottles in minimal essential medium (alpha modification) with 10% fetal bovine serum. The gene sequence of the human NK-2 receptor is given in N.P. Gerard, et al., Journal of Biological Chemistry, 265:20455-20462 (1990).

For preparation of membranes, 30 confluent roller bottle cultures were dissociated by washing each roller bottle with 10 ml of Dulbecco's phosphate buffered saline (PBS) without calcium and magnesium, followed by addition of 10 ml of enzyme-free cell dissociation solution (PBS-based, from Specialty Media, Inc.). After an additional 15 minutes, the dissociated cells were pooled and centrifuged at 1,000 RPM for 10 minutes in a clinical centrifuge. Membranes were prepared by homogenization of the cell pellets in 300 ml 50 mM Tris buffer, pH 7.4 with a Tekmar® homogenizer for 10-15 seconds, followed by

- 68 -

centrifugation at 12,000 RPM (20,000 x g) for 30 minutes using a Beckman JA-14® rotor. The pellets were washed once using the above procedure. and the final pellets were resuspended in 100-120 ml 50 mM Tris buffer, pH 7.4, and 4 ml aliquots stored frozen at -70°C. The protein
5 concentration of this preparation was 2 mg/ml.

For the receptor binding assay, one 4-ml aliquot of the CHO-hNK-2R membrane preparation was suspended in 40 ml of assay buffer containing 50 mM Tris, pH 7.4, 3 mM manganese chloride, 0.02% bovine serum albumin (BSA) and 4 µg/ml chymostatin. A 200 µl volume
10 of the homogenate (40 µg protein) was used per sample. The radioactive ligand was [¹²⁵I]iodohistidyl-neurokinin A (New England Nuclear, NEX-252), 2200 Ci/mmol. The ligand was prepared in assay buffer at 20 nCi per 100 µl; the final concentration in the assay was 20 pM. Non-specific binding was determined using 1 µM eledoisin. Ten
15 concentrations of eledoisin from 0.1 to 1000 nM were used for a standard concentration-response curve.

All samples and standards were added to the incubation in 10 µl dimethylsulfoxide (DMSO) for screening (single dose) or in 5 µl DMSO for IC₅₀ determinations. The order of additions for incubation was
20 190 or 195 µl assay buffer, 200 µl homogenate, 10 or 5 µl sample in DMSO, 100 µl radioactive ligand. The samples were incubated 1 hr at room temperature and then filtered on a cell harvester through filters which had been presoaked for two hours in 50 mM Tris buffer, pH 7.7, containing 0.5% BSA. The filter was washed 3 times with approximately
25 3 ml of cold 50 mM Tris buffer, pH 7.7. The filter circles were then punched into 12 x 75 mm polystyrene tubes and counted in a gamma counter.

Animal and human clinical models demonstrating the
30 effectiveness of the methods of the present invention are well known to those skilled in the art. For example, the following experiment clearly demonstrates the inhibitory effect of the compounds of the present invention on an animal model predictive of migraine therapies.

- 69 -

Neurogenic Plasma Estravasation in the Dural Layer Induced by
Electrical Stimulation

5 Harlan Sprague-Dawley rats (225-325 g) or guinea pigs from
Charles River Laboratories (225-325 g) are anesthetized with sodium
phenobarbitol (65 mg/kg or 45 mg/kg, respectively, intraperitoneally) and
placed in a stereotaxic frame (David Kopf Instruments) with the incisor
10 bar set at -3.5 mm for rats or -4.0 mm for guinea pigs. Following a
midline sagittal scalp incision, two pairs of bilateral holes are drilled
through the skull (6 mm posteriorly, 2.0 and 4.0 mm laterally for rats; 4
mm posteriorly and 3.2 and 5.2 mm laterally for guinea pigs -- all
coordinates reference to bregma). Pairs of stainless steel stimulating
electrodes, insulated except for the tips, are lowered through the holes in
15 both hemispheres to a depth of 9 mm (rats) or 10.5 mm (guinea pigs) from
dura.

 The femoral vein is exposed and a dose of the test compound
is injected intravenously (1 ml/kg). Approximately seven minutes later, a
50 mg/kg dose of Evans Blue, a fluorescent dye, is also injected
20 intravenously. The Evans Blue complexed with proteins in the blood and
functioned as a marker for protein extravasation. Exactly ten minutes
post-injection of the test compound, the left trigeminal ganglion is
stimulated for three minutes at a current intensity of 1.0 mA (5 Hz, 4
msec duration) with a potentiostat/galvanostat.

25 Fifteen minutes following the stimulation, the animals are
killed and exanguinated with 20 ml of saline. The top of the skull is
removed to facilitate the collection of the dural membranes. The
membrane samples are removed from both hemispheres, rinsed with
water, and spread flat on microscopic slides. Once dried, the tissues are
30 coverslipped with a 70% glycerol/water solution.

 A fluorescence microscope equipped with a grating
monochromator and a spectrophotometer is used to quantify the amount
of Evans Blue dye in each tissue sample. An excitation wavelength of

- 70 -

approximately 535 nm is utilized and the emission intensity at 600 nm is determined. The microscope is equipped with a motorized stage and is interfaced with a personal computer. This facilitated the computer-controlled movement of the stage with fluorescence measurements at 25
5 points (500 μ m steps) on each dural sample. The mean and standard deviation of the measurements are determined by the computer.

The dural extravasation induced by electrical stimulation of the trigeminal ganglion is an ipsilateral effect (i.e. it occurs only on the side of the dura in which the trigeminal ganglion is stimulated). This
10 allowed the other, unstimulated, half of the dura to be used as a control. The ratio of the amount of extravasation in the dura from the stimulated side compared to the unstimulated side is calculated. Saline controls yielded a ratio of approximately 2.0 in rats and 1.8 in guinea pigs. In contrast, a compound which effectively prevented the extravasation in the
15 dura from the stimulated side would have a ratio of approximately 1.0. A dose-response curve is generated and the dose that inhibited the extravasation by 50% (ID₅₀) is estimated.

The compounds prepared by the processes of the present
20 invention are useful as tachykinin receptor-binding compounds. As such, they may be employed as antagonists or agonists of the various tachykinins. These compounds are, therefore, useful in the treatment or prevention of conditions associated with an excess or deficiency of tachykinins. The term "physiological disorder associated with an excess
25 or deficiency of tachykinins" encompasses those disorders associated with an inappropriate stimulation of tachykinin receptors, regardless of the actual amount of tachykinin present in the locale.

These physiological disorders may include disorders of the central nervous system such as anxiety, depression, psychosis, and
30 schizophrenia; neurodegenerative disorders such as dementia, including senile dementia of the Alzheimer's type, Alzheimer's disease, AIDS-associated dementia, and Down's syndrome; demyelinating diseases such as multiple sclerosis and amyotrophic lateral sclerosis and other

- 71 -

neuropathological disorders such as peripheral neuropathy, such as diabetic and chemotherapy-induced neuropathy, and post-herpetic and other neuralgias; acute and chronic obstructive airway diseases such as adult respiratory distress syndrome, bronchopneumonia, bronchospasm, chronic bronchitis, drivercough, and asthma; inflammatory diseases such as inflammatory bowel disease, psoriasis, fibrositis, osteoarthritis, and rheumatoid arthritis; disorders of the musculo-skeletal system, such as osteoporosis; allergies such as eczema and rhinitis; hypersensitivity disorders such as poison ivy; ophthalmic diseases such as conjunctivitis, vernal conjunctivitis, and the like; cutaneous diseases such as contact dermatitis, atopic dermatitis, urticaria, and other eczematoid dermatites; addiction disorders such as alcoholism; stress-related somatic disorders; reflex sympathetic dystrophy such as shoulder/hand syndrome; dysthymic disorders; adverse immunological reactions such as rejection of transplanted tissues and disorders related to immune enhancement or suppression such as systemic lupus erythematosus; gastrointestinal disorders or diseases associated with the neuronal control of viscera such as ulcerative colitis, Crohn's disease and irritable bowel syndrome; disorders of bladder function such as bladder detrusor hyper-reflexia and incontinence; atherosclerosis; fibrosing and collagen diseases such as scleroderma and eosinophilic fascioliasis; irritative symptoms of benign prostatic hypertrophy; disorders of blood flow caused by vasodilation and vasospastic diseases such as angina, migraine, and Reynaud's disease; emesis; and pain or nociception, for example, that attributable to or associated with any of the foregoing conditions, especially the transmission of pain in migraine. For example the compounds of Formula I may suitably be used in the treatment of disorders of the central nervous system such as anxiety, psychosis, and schizophrenia; neurodegenerative disorders such as Alzheimer's disease and Down's syndrome; respiratory diseases such as bronchospasm and asthma; inflammatory diseases such as inflammatory bowel disease, osteoarthritis and rheumatoid arthritis; adverse immunological disorders such as rejection of transplanted tissues; gastrointestinal disorders and diseases such as disorders associated with

- 72 -

the neuronal control of viscera such as ulcerative colitis, Crohn's disease and irritable bowel syndrome; incontinence; disorders of blood flow caused by vasodilation; and pain or nociception, for example, that attributable to or associated with any of the foregoing conditions or the transmission of
5 pain in migraine.

The results of several experiments demonstrate that many of the compounds of Formula I are selective tachykinin receptor antagonists. These compounds preferentially bind one tachykinin receptor subtype compared to other such receptors. Such compounds are especially
10 preferred.

For example, NK-1 antagonists are most especially preferred in the treatment of pain, especially chronic pain, such as neuropathic pain, post-operative pain, and migraines, pain associated with arthritis, cancer-associated pain, chronic lower back pain, cluster headaches,
15 herpes neuralgia, phantom limb pain, central pain, dental pain, sunburn pain, neuropathic pain, opioid-resistant pain, visceral pain, surgical pain, bone injury pain, pain during labor and delivery, pain resulting from burns, post partum pain, angina pain, and genitourinary tract-related pain including cystitis.

In addition to pain, NK-1 antagonists are especially
preferred in the treatment and prevention of urinary incontinence;
irritative symptoms of benign prostatic hypertrophy; motility disorders of
the gastrointestinal tract, such as irritable bowel syndrome; acute and chronic obstructive airway diseases, such as bronchospasm,
20 bronchopneumonia, asthma, and adult respiratory distress syndrome; atherosclerosis; inflammatory conditions, such as inflammatory bowel disease, ulcerative colitis, Crohn's disease, rheumatoid arthritis, osteoarthritis, neurogenic inflammation, allergies, rhinitis, cough, dermatitis, urticaria, psoriasis, conjunctivitis, irritation-induced miosis;
25 tissue transplant rejection; plasma extravasation resulting from cytokine chemotherapy and the like; spinal cord trauma; stroke; cerebral stroke (ischemia); Alzheimer's disease; Parkinson's disease; multiple sclerosis; amyotrophic lateral sclerosis; schizophrenia; anxiety; and depression.

- 73 -

NK-2 antagonists are especially preferred in the treatment of urinary incontinence, bronchospasm, asthma, adult respiratory distress syndrome, motility disorders of the gastrointestinal tract, such as irritable bowel syndrome, and pain.

5 In addition to the in vitro binding assays described supra, many of the compounds prepared by the processes of the present invention have also been tested in in vivo model systems for conditions associated with an excess of tachykinins. Of those compounds tested in vivo many have shown efficacy against said conditions.

10 While it is possible to administer a compound employed in the methods of this invention directly without any formulation, the compounds are usually administered in the form of pharmaceutical compositions comprising a pharmaceutically acceptable excipient and at
15 least one active ingredient. These compositions can be administered by a variety of routes including oral, rectal, transdermal, subcutaneous, intravenous, intramuscular, and intranasal. Many of the compounds employed in the methods of this invention are effective as both injectable and oral compositions. Such compositions are prepared in a manner well
20 known in the pharmaceutical art and comprise at least one active compound. See, e.g., REMINGTON'S PHARMACEUTICAL SCIENCES, (16th ed. 1980).

 In making the compositions employed in the present invention the active ingredient is usually mixed with an excipient, diluted
25 by an excipient or enclosed within such a carrier which can be in the form of a capsule, sachet, paper or other container. When the excipient serves as a diluent, it can be a solid, semi-solid, or liquid material, which acts as a vehicle, carrier or medium for the active ingredient. Thus, the compositions can be in the form of tablets, pills, powders, lozenges,
30 sachets, cachets, elixirs, suspensions, emulsions, solutions, syrups, aerosols (as a solid or in a liquid medium), ointments containing for example up to 10% by weight of the active compound, soft and hard

- 74 -

gelatin capsules, suppositories, sterile injectable solutions, and sterile packaged powders.

5 In preparing a formulation, it may be necessary to mill the active compound to provide the appropriate particle size prior to combining with the other ingredients. If the active compound is substantially insoluble, it ordinarily is milled to a particle size of less than 200 mesh. If the active compound is substantially water soluble, the particle size is normally adjusted by milling to provide a substantially uniform distribution in the formulation, e.g. about 40 mesh.

10 Some examples of suitable excipients include lactose, dextrose, sucrose, sorbitol, mannitol, starches, gum acacia, calcium phosphate, alginates, tragacanth, gelatin, calcium silicate, microcrystalline cellulose, polyvinylpyrrolidone, cellulose, water, syrup, and methyl cellulose. The formulations can additionally include:
15 lubricating agents such as talc, magnesium stearate, and mineral oil; wetting agents; emulsifying and suspending agents; preserving agents such as methyl- and propylhydroxybenzoates; sweetening agents; and flavoring agents. The compositions of the invention can be formulated so as to provide quick, sustained or delayed release of the active ingredient
20 after administration to the patient by employing procedures known in the art.

The compositions are preferably formulated in a unit dosage form, each dosage containing from about 0.05 to about 100 mg, more usually about 1.0 to about 30 mg, of the active ingredient. The term "unit
25 dosage form" refers to physically discrete units suitable as unitary dosages for human subjects and other mammals, each unit containing a predetermined quantity of active material calculated to produce the desired therapeutic effect, in association with a suitable pharmaceutical excipient.

30 The active compounds are generally effective over a wide dosage range. For examples, dosages per day normally fall within the range of about 0.01 to about 30 mg/kg of body weight. In the treatment of adult humans, the range of about 0.1 to about 15 mg/kg/day, in single or

- 75 -

divided dose, is especially preferred. However, it will be understood that the amount of the compound actually administered will be determined by a physician, in the light of the relevant circumstances, including the condition to be treated, the chosen route of administration, the actual
5 compound or compounds administered, the age, weight, and response of the individual patient, and the severity of the patient's symptoms, and therefore the above dosage ranges are not intended to limit the scope of the invention in any way. In some instances dosage levels below the
10 lower limit of the aforesaid range may be more than adequate, while in other cases still larger doses may be employed without causing any harmful side effect, provided that such larger doses are first divided into several smaller doses for administration throughout the day.

- 76 -

Formulation Preparation 1

Hard gelatin capsules containing the following ingredients
are prepared:

5

<u>Ingredient</u>		<u>Quantity</u>
Active Ingredient		(mg/capsule)
		30.0
10	Starch	305.0
	Magnesium stearate	5.0

The above ingredients are mixed and filled into hard gelatin
15 capsules in 340 mg quantities.

- 77 -

Formulation Preparation 2

A tablet formula is prepared using the ingredients below:

5		Quantity
	<u>Ingredient</u>	<u>(mg/tablet)</u>
	Active Ingredient	25.0
	Cellulose, microcrystalline	200.0
10	Colloidal silicon dioxide	10.0
	Stearic acid	5.0

15 The components are blended and compressed to form tablets,
each weighing 240 mg.

Formulation Preparation 3

20 A dry powder inhaler formulation is prepared containing the
following components:

	<u>Ingredient</u>	<u>Weight %</u>
	Active Ingredient	5
25	Lactose	95

 The active mixture is mixed with the lactose and the mixture
is added to a dry powder inhaling appliance.

30

- 78 -

Formulation Preparation 4

Tablets, each containing 30 mg of active ingredient, are prepared as follows:

5

<u>Ingredient</u>		<u>Quantity</u> <u>(mg/tablet)</u>
Active Ingredient		30.0 mg
10	Starch	45.0 mg
	Microcrystalline cellulose	35.0 mg
15	Polyvinylpyrrolidone (as 10% solution in water)	4.0 mg
	Sodium carboxymethyl starch	4.5 mg
	Magnesium stearate	0.5 mg
20	Talc	<u>1.0 mg</u>
	Total	120 mg

25

The active ingredient, starch and cellulose are passed through a No. 20 mesh U.S. sieve and mixed thoroughly. The solution of polyvinylpyrrolidone is mixed with the resultant powders, which are then passed through a 16 mesh U.S. sieve. The granules so produced are dried at 50-60°C and passed through a 16 mesh U.S. sieve. The sodium

30 carboxymethyl starch, magnesium stearate, and talc, previously passed through a No. 30 mesh U.S. sieve, are then added to the granules which, after mixing, are compressed on a tablet machine to yield tablets each weighing 120 mg.

- 79 -

Formulation Preparation 5

Capsules, each containing 40 mg of medicament are made as follows:

5

<u>Ingredient</u>	<u>Quantity</u> <u>(mg/capsule)</u>
Active Ingredient	40.0 mg
Starch	109.0 mg
Magnesium stearate	<u>1.0 mg</u>
Total	150.0 mg

15

The active ingredient, cellulose, starch, and magnesium stearate are blended, passed through a No. 20 mesh U.S. sieve, and filled into hard gelatin capsules in 150 mg quantities.

20

Formulation Preparation 6

Suppositories, each containing 25 mg of active ingredient are made as follows:

25

<u>Ingredient</u>	<u>Amount</u>
Active Ingredient	25 mg
Saturated fatty acid glycerides to	2,000 mg

30

The active ingredient is passed through a No. 60 mesh U.S. sieve and suspended in the saturated fatty acid glycerides previously melted using the minimum heat necessary. The mixture is then poured into a suppository mold of nominal 2.0 g capacity and allowed to cool.

- 80 -

Formulation Preparation 7

Suspensions, each containing 50 mg of medicament per 5.0 ml dose are made as follows:

	<u>Ingredient</u>	<u>Amount</u>
	Active Ingredient	50.0 mg
10	Xanthan gum	4.0 mg
	Sodium carboxymethyl cellulose (11%)	
	Microcrystalline cellulose (89%)	50.0 mg
15	Sucrose	1.75 g
	Sodium benzoate	10.0 mg
	Flavor and Color	q.v.
20	Purified water to	5.0 ml

The medicament, sucrose and xanthan gum are blended, passed through a No. 10 mesh U.S. sieve, and then mixed with a previously made solution of the microcrystalline cellulose and sodium carboxymethyl cellulose in water. The sodium benzoate, flavor, and color are diluted with some of the water and added with stirring. Sufficient water is then added to produce the required volume.

- 81 -

Formulation Preparation 8

5 Capsules, each containing 15 mg of medicament, are made as follows:

	<u>Ingredient</u>	<u>Quantity</u> <u>(mg/capsule)</u>
	Active Ingredient	15.0 mg
10	Starch	407.0 mg
	Magnesium stearate	<u>3.0 mg</u>
15	Total	425.0 mg

20 The active ingredient, cellulose, starch, and magnesium stearate are blended, passed through a No. 20 mesh U.S. sieve, and filled into hard gelatin capsules in 425 mg quantities.

Formulation Preparation 9

An intravenous formulation may be prepared as follows:

	<u>Ingredient</u>	<u>Quantity</u>
25	Active Ingredient	250.0 mg
	Isotonic saline	1000 ml
30		

- 82 -

Formulation Preparation 10

A topical formulation may be prepared as follows:

5	<u>Ingredient</u>	<u>Quantity</u>
	Active Ingredient	1-10 g
	Emulsifying Wax	30 g
10	Liquid Paraffin	20 g
	White Soft Paraffin	to 100 g

- 15 The white soft paraffin is heated until molten. The liquid paraffin and emulsifying wax are incorporated and stirred until dissolved. The active ingredient is added and stirring is continued until dispersed. The mixture is then cooled until solid.

- 83 -

Formulation Preparation 11

Sublingual or buccal tablets, each containing 10 mg of active ingredient, may be prepared as follows:

5	<u>Ingredient</u> Active Ingredient	<u>Quantity</u> <u>Per Tablet</u> 10.0 mg
10	Glycerol	210.5 mg
	Water	143.0 mg
	Sodium Citrate	4.5 mg
15	Polyvinyl Alcohol	26.5 mg
	Polyvinylpyrrolidone	<u>15.5 mg</u>
	Total	410.0 mg

20 The glycerol, water, sodium citrate, polyvinyl alcohol, and polyvinylpyrrolidone are admixed together by continuous stirring and maintaining the temperature at about 90°C. When the polymers have gone into solution, the solution is cooled to about 50-55°C and the

25 medicament is slowly admixed. The homogenous mixture is poured into forms made of an inert material to produce a drug-containing diffusion matrix having a thickness of about 2-4 mm. This diffusion matrix is then cut to form individual tablets having the appropriate size.

30 Another preferred formulation employed in the methods of the present invention employs transdermal delivery devices ("patches"). Such transdermal patches may be used to provide continuous or discontinuous infusion of the compounds of the present invention in

- 84 -

controlled amounts. The construction and use of transdermal patches for the delivery of pharmaceutical agents is well known in the art. See, e.g., U.S. Patent 5,023,252, issued June 11, 1991, herein incorporated by reference. Such patches may be constructed for continuous, pulsatile, or on demand delivery of pharmaceutical agents.

Frequently, it will be desirable or necessary to introduce the pharmaceutical composition to the brain, either directly or indirectly. Direct techniques usually involve placement of a drug delivery catheter into the host's ventricular system to bypass the blood-brain barrier. One such implantable delivery system, used for the transport of biological factors to specific anatomical regions of the body, is described in U.S. Patent 5,011,472, issued April 30, 1991, which is herein incorporated by reference.


Indirect techniques, which are generally preferred, usually involve formulating the compositions to provide for drug latentiation by the conversion of hydrophilic drugs into lipid-soluble drugs or prodrugs. Latentiation is generally achieved through blocking of the hydroxy, carbonyl, sulfate, and primary amine groups present on the drug to render the drug more lipid soluble and amenable to transportation across the blood-brain barrier. Alternatively, the delivery of hydrophilic drugs may be enhanced by intra-arterial infusion of hypertonic solutions which can transiently open the blood-brain barrier.

The type of formulation employed for the administration of the compounds employed in the methods of the present invention may be dictated by the particular compounds employed, the type of pharmacokinetic profile desired from the route of administration and the compound, and the state of the patient.

- 85 -

We Claim:

1. A compound of the formula



5

wherein:

10 R^1 and R^2 are independently hydrogen, halo, C_1 - C_6 alkyl, hydroxy, or C_1 - C_6 alkoxy;

R^5 , R^6 , and R^7 , are independently hydrogen, halo, C_1 - C_6 alkyl, C_1 - C_6 alkoxy, trifluoromethyl, or hydroxy;

15 R^3 is hydrogen, C_2 - C_7 alkanoyl, glycyl, or dimethylglycyl;

n is 1-6;

D is $-S(O)_m-$, $-NH-$, $-C(O)-$, or $-O-$,

20

m is 0, 1, or 2;

- 86 -

5 R⁸ and R⁹ are independently hydrogen, hydroxy, C₁-C₆ alkyl, C₁-C₆ alkoxy, C₁-C₆ alkylthio, C₁-C₆ alkoxy(C₁-C₆ alkylenyl)-, C₂-C₇ alkoxycarbonyl, C₂-C₇ alkoxycarbonyl(C₁-C₆ alkylenyl)-, trifluoromethoxy, trichloromethoxy, C₁-C₆ alkylthio, C₁-C₆ alkylamino, di(C₁-C₆ alkyl)amino, formyl, cyano, halo, trifluoromethyl, R¹⁰R¹¹N(C₁-C₆ alkylenyl)-, pyrrolyl, triazolyl, imidazolyl, tetrazolyl, thiazolyl, thiazolinyl, thiadiazolyl, thiadiazolinyl, piperidyl, pyrrolidyl, morpholinyl, morpholinocarbonyl, hexamethyleneiminyl, methylsulfonyl, methylsulfinyl, phenoxy, benzyloxy, carboxy, carbamoyl, or C₂-C₇ alkylcarbamoyl(C₁-C₆ alkylenyl)-,

15 R¹⁰ and R¹¹ are independently hydrogen or C₁-C₆ alkyl,


said C₁-C₆ alkyl or C₁-C₆ alkoxy groups being optionally substituted with one, two, or three moieties selected from the group consisting of hydroxy, halo, cyano, amino, nitro, carboxy, carbamoyl, and thiol;

25 or R⁸ and R⁹ can combine, together with the benzo ring to which they are attached, to form a naphthyl, dihydronaphthyl, tetrahydronaphthyl, quinolinyl, isoquinolinyl, 2-coumaranonyl, 3-coumaranonyl, benzothiazolyl, benzimidazolyl, indolyl, benzothienyl, benzofuryl, 2,3-dihydrobenzofuryl, indolinyl, or 2,3-dihydrobenzothienyl group, which can be attached to D at
30 any position of the bicyclic group;

or a pharmaceutically acceptable salt or solvate thereof.

- 87 -

2. A method of treating a condition associated with an excess of tachykinins, which comprises administering to a mammal in need thereof an effective amount of a compound of the formula



wherein:

10

R^1 and R^2 are independently hydrogen, halo, C_1 - C_6 alkyl, hydroxy, or C_1 - C_6 alkoxy;

15

R^5 , R^6 , and R^7 , are independently hydrogen, halo, C_1 - C_6 alkyl, C_1 - C_6 alkoxy, trifluoromethyl, or hydroxy;

R^3 is hydrogen, C_2 - C_7 alkanoyl, glycyl, or dimethylglycyl;

n is 1-6;

20

D is $-S(O)_m-$, $-NH-$, $-C(O)-$, or $-O-$,

m is 0, 1, or 2;

- 88 -


- 5 R⁸ and R⁹ are independently hydrogen, hydroxy, C₁-C₆ alkyl, C₁-C₆ alkoxy, C₁-C₆ alkylthio, C₁-C₆ alkoxy(C₁-C₆ alkylenyl)-, C₂-C₇ alkoxycarbonyl, C₂-C₇ alkoxycarbonyl(C₁-C₆ alkylenyl)-, trifluoromethoxy, trichloromethoxy, C₁-C₆ alkylthio, C₁-C₆ alkylamino, di(C₁-C₆ alkyl)amino, formyl, cyano, halo, trifluoromethyl, R¹⁰R¹¹N(C₁-C₆ alkylenyl)-, pyrrolyl, triazolyl, imidazolyl, tetrazolyl, thiazolyl, thiazolinyl, thiadiazolyl, thiadiazolinyl, piperidyl, pyrrolidyl, morpholinyl, morpholinocarbonyl, hexamethyleneiminy, methylsulfonyl, methylsulfinyl, phenoxy, benzyloxy, carboxy, carbamoyl, or C₂-C₇ alkylcarbamoyl(C₁-C₆ alkylenyl)-,
- 10
- 15 R¹⁰ and R¹¹ are independently hydrogen or C₁-C₆ alkyl,
- 20 said C₁-C₆ alkyl or C₁-C₆ alkoxy groups being optionally substituted with one, two, or three moieties selected from the group consisting of hydroxy, halo, cyano, amino, nitro, carboxy, carbamoyl, and thiol;
- 25 or R⁸ and R⁹ can combine, together with the benzo ring to which they are attached, to form a naphthyl, dihydronaphthyl, tetrahydronaphthyl, quinolinyl, isoquinolinyl, 2-coumaranonyl, 3-coumaranonyl, benzothiazolyl, benzimidazolyl, indolyl, benzothienyl, benzofuryl, 2,3-dihydrobenzofuryl, indolinyl, or 2,3-dihydrobenzothienyl group, which can be attached to D at
- 30 any position of the bicyclic group;

- 89 -

or a pharmaceutically acceptable salt or solvate thereof.

3. A pharmaceutical formulation, comprising a compound of the formula

5



wherein:

10 R^1 and R^2 are independently hydrogen, halo, C_1 - C_6 alkyl, hydroxy, or C_1 - C_6 alkoxy;

R^5 , R^6 , and R^7 , are independently hydrogen, halo, C_1 - C_6 alkyl, C_1 - C_6 alkoxy, trifluoromethyl, or hydroxy;

15

R^3 is hydrogen, C_2 - C_7 alkanoyl, glycyl, or dimethylglycyl;

n is 1-6;

20

D is $-S(O)_m-$, $-NH-$, $-C(O)-$, or $-O-$,

m is 0, 1, or 2;

- 90 -

- 5 R⁸ and R⁹ are independently hydrogen, hydroxy, C₁-C₆ alkyl, C₁-C₆ alkoxy, C₁-C₆ alkylthio, C₁-C₆ alkoxy(C₁-C₆ alkylenyl)-, C₂-C₇ alkoxycarbonyl, C₂-C₇ alkoxycarbonyl(C₁-C₆ alkylenyl)-, trifluoromethoxy, trichloromethoxy, C₁-C₆ alkylthio, C₁-C₆ alkylamino, di(C₁-C₆ alkyl)amino, formyl, cyano, halo, trifluoromethyl, R¹⁰R¹¹N(C₁-C₆ alkylenyl)-, pyrrolyl, triazolyl, imidazolyl, tetrazolyl, thiazolyl, thiazolinyl, thiadiazolyl, thiadiazolinyl, piperidyl, pyrrolidyl, morpholinyl, morpholinocarbonyl, hexamethyleneiminyl, methylsulfonyl, methylsulfinyl, phenoxy, benzyloxy, carboxy, carbamoyl, or C₂-C₇ alkylcarbamoyl(C₁-C₆ alkylenyl)-,
- 10
- 15 R¹⁰ and R¹¹ are independently hydrogen or C₁-C₆ alkyl,
- said C₁-C₆ alkyl or C₁-C₆ alkoxy groups being optionally substituted with one, two, or three
- 20 moieties selected from the group consisting of hydroxy, halo, cyano, amino, nitro, carboxy, carbamoyl, and thiol;
- or R⁸ and R⁹ can combine, together with the benzo ring to which they are attached, to form a naphthyl, dihydronaphthyl, tetrahydronaphthyl, quinolinyl, isoquinolinyl, 2-coumaranonyl, 3-coumaranonyl, benzothiazolyl, benzimidazolyl, indolyl, benzothienyl, benzofuryl, 2,3-dihydrobenzofuryl, indolinyl, or 2,3-
- 25
- 30 dihydrobenzothienyl group, which can be attached to D at any position of the bicyclic group;

- 91 -

or a pharmaceutically acceptable salt or solvate thereof, in combination with one or more pharmaceutically acceptable carriers, diluents, or excipients therefor.

INTERNATIONAL SEARCH REPORT

International application No.
PCT/US98/16333

A. CLASSIFICATION OF SUBJECT MATTER

IPC(6) :C07D 209/16; A01N 43/38; A61K 31/40, 31/405

US CL :514/412,415,419; 548/455,504,506

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

U.S. : 514/412,415,419; 548/455,504,506

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practicable, search terms used)

STN structure search in Registry and Chemical Abstracts files

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	Chemical Abstract, Volume 122, Abstract No. 122:32016, HORWELL et al, "Preparation of N-substituted cycloalkyl and polycycloalkyl alpha-substituted tryptophanylphenylalanine derivatives as drugs", abstract for US 5,278,316 A, see the entire abstract.	1-3



Further documents are listed in the continuation of Box C.



See patent family annex.

* Special categories of cited documents:	*T* later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
A document defining the general state of the art which is not considered to be of particular relevance	*X* document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
B earlier document published on or after the international filing date	*Y* document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art
L document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)	*Z* document member of the same patent family
O document referring to an oral disclosure, use, exhibition or other means	
P document published prior to the international filing date but later than the priority date claimed	

Date of the actual completion of the international search

10 SEPTEMBER 1998

Date of mailing of the international search report

13 OCT 1998

Name and mailing address of the ISA/US
Commissioner of Patents and Trademarks
Box PCT
Washington, D.C. 20231

Facsimile No. (703) 305-3230

Authorized officer

FREDERICK KRASS

Telephone No. (703) 308-2351

14. A method of screening for a compound that modulates activity of T-type calcium channel α_{11} subunit variants TCCV-1 and TCCV-2 having the amino acid sequence of SEQ ID NO: 2 and 4 respectively, the method comprising:

- (a) contacting a TCCV-1 or TCCV-2 polypeptide with the compound; and
- (b) detecting modulation of activity of the TCCV-1 or TCCV-2 polypeptide.

14. A method of screening for a compound that modulates activity of T-type calcium channel α_{1I} subunit variants TCCV-1 and TCCV-2 having the amino acid sequences of SEQ ID NO: 2 and 4 respectively, the method comprising:

- (a) contacting a TCCV-1 or TCCV-2 polypeptide with the compound; and
- (b) detecting modulation of activity of the TCCV-1 or TCCV-2 polypeptide.

Material Transfer Agreement

This Material Transfer Agreement ("MTA") is entered into as of this 27th day of January, 2003, ("Effective Date") between Roche Palo Alto LLC having offices at 3401 Hillview Avenue, Palo Alto CA 94304 and Genetastix Corporation ("Genetastix") having an office at 780 Montague Expressway, Suite 303, San Jose, CA 95131.

1. **Material to be Transferred.** Genetastix will transfer certain protein (or proteins), which are antihuman CCR5 scFv antibodies related to certain expression technology and whose manufacture is proprietary to Genetastix ("Materials"). "Materials" also includes, in addition to the original biological and/or other materials transferred to Roche, any derivatives, progeny, or improvements developed by Roche from those materials, and any combination of the foregoing with other substances. Upon execution of this Agreement, the parties will make arrangements for delivery of one or more samples of Materials within five (5) days of the Effective Date, or with arrangements that both parties agree
2. **Purpose of Transfer; Use.** The Materials may be used solely for conducting the experiments described in Exhibit A (the "Experiments") by Roche at Roche' facilities, in order for Genetastix and Roche to evaluate whether they wish to enter into a further transaction related to the Materials. The Materials may not be sold or used for any other purposes, including commercial or production purposes. Roche may not reverse engineer, modify, extract other materials from, or develop any products incorporating, based on or derived from the Materials. Without limiting the foregoing, Roche may not make any modifications (whether by mutagenesis, mutations, deletions, insertions or otherwise) with respect to the variable heavy, light chain and the constant region in the Materials.
3. **Results.** Roche will communicate to Genetastix in writing, within thirty (30) days after completion of the Experiments, a summary of the arising from the Experiments ("Results").
4. **Confidentiality and Use of Information and Results.** Information associated with the Materials and made available to Roche under this Agreement, whether directly or indirectly by Genetastix ("Information") is and will remain proprietary and confidential to Genetastix. Roche agrees to hold all Information and Results in confidence, to restrict access to the Information and Results as required under this Agreement, not to disclose the Information or Results to any third party, other than its Affiliates and not to use the Information or Results for any purpose except the conduct of the Experiments and the related evaluation. Roche shall not be subject to the obligations set forth above as to the disclosure or use of any Information that: (1) is or later becomes publicly known under circumstances involving no breach of this Agreement by Roche; (2) is already known to Roche or its Affiliates at the time of receipt; (3) is lawfully made available to Roche or its Affiliates by a third party, in each case, as evidenced by Roche's or its Affiliate's written records kept in the ordinary course of business, or (4) is generated at Roche or its Affiliates by persons who have not had access to or knowledge of the Information disclosed hereunder.

5. **Care and Access.** Roche will use prudence and reasonable care in the use, handling, storage, transportation and disposition of the Materials, and shall use the Materials in compliance with all applicable national, state and local laws and regulations, including without limitation all applicable National Institutes of Health Guidelines.
6. **Payments to Genetastix.** Roche will make a non-refundable, non-creditable payment to Genetastix of \$25,000 US within five (5) business days of receipt of Materials under this MTA.
7. **No License Granted.** Roche understands that the supply of the Materials shall not be construed as granting any license, grant or right under any of Genetastix's patents or other rights other than to carry out the experiments proposed herein.
8. **Limitations on Use.** Without limiting the other restrictions regarding the Materials, Information or Results, Roche shall not use the Materials, Information, or Results or any derivatives or portions of the foregoing, in support of any of its any patent applications.
9. **Term of MTA; Negotiation for Potential License.** The term of this MTA is one hundred and twenty (120) days from the Effective Date, during which the parties may begin negotiations for a license agreement. If the parties do not reach such an agreement within ninety (90) days of beginning negotiations, Roche shall discontinue any use of the Materials, Information or Results and will destroy any remaining Material.
10. **Warranty.** Both parties further warrant that as of the Effective Date the terms and conditions of this MTA are not inconsistent with any other agreements or obligations that the parties may have. Other than as set forth herein, Roche agrees that Genetastix makes no warranties whatsoever regarding the quality of the Materials. GENETASTIX DISCLAIMS ANY WARRANTIES OF MERCHANTABILITY, FITNESS FOR A PARTICULAR PURPOSE, AND ANY OTHER WARRANTY OTHER THAN THOSE IN THIS SECTION, WHETHER EXPRESS OR IMPLIED, AND ARISING OUT OF STATUTE, COURSE OF DEALING OR OTHERWISE.
11. **Third Party Information.** Genetastix does not wish to receive information Roche considers to be confidential and outside of the studies contemplated hereunder. Both parties will not disclose to each other confidential information of any third party.
12. **Hold Harmless.** Each party is responsible for its own acts and omissions relating to the Experiments, the Materials, and its conduct under this MTA. The parties each will indemnify and hold the other harmless, along with their employees, officers, agents, directors, and contractors, from and against any and all loss and/or liability (including but not limited to attorney's fees and other related costs) arising out of personal injury or death of its employees, agents, on contractors, or loss of or

damage to its tangible personal property, in connection with its performance under this MTA, except to the extent that such injury, death, loss or damage is determined, after final judgment from which no appeal can be taken, to have resulted from an act or omission amounting to sole negligence or willful misconduct of the other party or an employee, agent, officer, or representative thereof.

- 13. Disputes.** This MTA will be governed by, and construed and interpreted in accordance with, the laws of the state of California, without reference to conflicts of laws principles. Any claim or controversy arising out of or related to this MTA, or any breach of this MTA, must be submitted to a court of applicable and competent jurisdiction in the state of California, and each party hereby consents to the jurisdiction and venue of such a court.
- 14. Assignment.** Roche shall not assign, or attempt to assign, this MTA, whether by operation of law, a change in control or otherwise, without the prior express written consent of Genetastix, and any purported assignment in violation of the foregoing is void and of no effect.

GENETASTIX CORPORATION ROCHE PALO ALTO LLC.

By: _____ By: _____

Name: _____ Name: _____

Title: _____ Title: _____

Contacting Party:

Li Zhu, Ph.D, President and CEO, Genetastix Corporation
780 Montague Expressway, Suite 303, San Jose, CA 95131
Phone: 408-435-8900 x101, Fax: 408-433-9288
E-mail: li.zhu@genetastix.com

Exhibit A

Experiments

Roche may:

1. Receptor ligand binding assays using hCCR5 receptor transfected CHO cells
2. FLIPR assay using hCCR5 transfected receptor CHO cells
3. Displacement of ¹²⁵I-gp120 binding from live CHO-hCCR5 cells
4. Inhibition of virus-induced infectious focus formation in MAGI-CCR5 cells